

03 JAN 2005 20, 153

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number
WO 2004/005498 A1

(51) International Patent Classification⁷: **C12N 7/01**,
15/63, C12Q 1/70, A61K 39/29

1517 Bayou Shore Dr., Galveston, TX 77551 (US). **RI-JNBRAND, Rene** [NL/US]; 1427 Winne, Galveston, TX 77550 (US).

(21) International Application Number:
PCT/US2003/021002

(74) Agent: **SHISHIMA, Gina, N.**; Fulbright & Jaworski L.L.P., 600 Congress Avenue, Suite 2400, Austin, TX 78701 (US).

(22) International Filing Date: 2 July 2003 (02.07.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
10/189,359 3 July 2002 (03.07.2002) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 10/189,359 (CON)
Filed on 3 July 2002 (03.07.2002)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants (*for all designated States except US*): **BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 West 7th St., Austin, TX 78701 (US). **INSTITUT PASTEUR** [FR/FR]; 25-28 rue du Docteur Roux, F-75724 Paris Cedex 15 (FR).

Published:
— with international search report

(72) Inventors; and
(75) Inventors/Applicants (*for US only*): **MARTIN, Annette** [FR/FR]; 201 rue d'Alesia, F-75014 Paris (FR). **SANGAR, David, V.** [GB/GB]; 23A Coventry Road, Burbage, Leicestershire LE10 2HL (GB). **LEMON, Stanley, M.** [US/US];

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CHIMERIC GB VIRUS B (GBV-B)

(57) Abstract: The present invention relates generally to the fields of biochemistry, molecular biology, and virology. More particularly, it relates to the identification of GB virus B (GBV-B)/HCV chimeras. The invention involves nucleic acid constructs and compositions encoding GBV-B/HCV chimera, including at least part of a 5' NTR derived from a HCV 5' NTR. This construct, and chimeric versions of it, may be employed to study GBV-B and related hepatitis family members, such as hepatitis C virus. The invention thus includes methods of preparing GBV-B/HCV chimeric sequences, constructs, and viruses, as well as methods of employing these compositions.

WO 2004/005498 A1

DESCRIPTION**CHIMERIC GB VIRUS B (GBV-B)****BACKGROUND OF THE INVENTION**

The present application claims priority to U.S. Patent Application Serial Number 10/189,359 filed on July 3, 2002. The entire text of the above-referenced disclosures is herein incorporated by reference.

A. Field of the Invention

The present invention relates generally to the fields of biochemistry, molecular biology, and virology. More particularly, it relates to the identification of 259 nucleotides of previously unrecognized sequence located at the 3' end of the GB virus B (GBV-B) genome.

B. Description of Related Art

Chronic hepatitis C is a major threat to the public health. Serologic surveys suggest that as many as 3.9 million Americans are chronically infected with the responsible virus, hepatitis C virus (HCV) (Alter, 1997). These individuals are at increased risk of developing progressive hepatic fibrosis leading to cirrhosis and loss of hepatocellular function, as well as hepatocellular carcinoma. The course of chronic hepatitis C is typically lengthy, often extending over decades, with insidious clinical progression usually occurring in the absence of symptoms. Nonetheless, liver disease due to HCV results in the death of 8,000-10,000 Americans annually, and chronic hepatitis C is the most common cause of liver transplantation within the U.S.

Therefore, HCV is a major public health problem. However, therapy for chronic hepatitis C is problematic. Recombinant interferon- α is approved for treatment of chronic hepatitis C (Consensus Development Panel, 1997). The benefit of interferon- α results primarily from its antiviral properties and its ability to inhibit production of virus by infected hepatocytes (Neumann *et al.*, 1998). Nonetheless, even under optimal therapeutic regimens, the majority of patients with chronic hepatitis C fail to eliminate the virus or resolve their liver disease. Treatment failures are especially common in persons infected with genotype 1 HCV, unfortunately the most prevalent genotype in the U.S. Thus, there is an urgent need to better understand the virus and develop better treatment. Unfortunately, technical difficulties in

working with HCV have made it necessary to use infectious surrogate viruses in efforts to develop treatments and vaccines for HCV.

Scientists' efforts to better understand HCV and to develop new drugs for treatment of hepatitis C have been stymied by two overwhelming technical deficiencies: first, the nonexistence of a high permissive cell line that supports replication of the virus and second, the absence of a permissive animal species other than chimpanzees, which are endangered and therefore available on a limited basis.

Presently, those who are working on HCV treatment and prevention are employing an infectious chimeric virus of sindbis and HCV and/or an infectious clone of pestiviruses as surrogate virus models in HCV drug discovery efforts, due to the above technical difficulties of working with HCV. Alternatively, they are using isolated proteins or RNA segments of HCV for biochemical and structural studies. This approach precludes functional studies of virus replication and its inhibition.

GBV-B is a hepatotropic flavivirus that has a unique phylogenetic relationship to human HCV and strong potential to serve as a surrogate virus in drug discovery efforts related to hepatitis C antiviral drug development. GBV-B causes acute hepatitis in experimentally infected tamarins (Simons *et al.*, 1995; Schlauder *et al.*, 1995; Karayiannis *et al.*, 1989) and can serve as a surrogate virus for HCV in drug discovery efforts (due to technical difficulties in working with HCV). GBV-B virus is much closer in sequence and biological properties than the above-described models. It will be easier to make biologically relevant chimeras between HCV and GBV-B than by using more distantly related viruses. GBV-B is hepatotropic (as is HCV), whereas the viruses used in these competing technologies are not. In view of the above, an infectious clone of GBV-B would be useful to those working on HCV treatment and prevention.

Unfortunately, the use of GBV-B as a surrogate or model for HCV has not been possible in the past, because no infectious molecular clone of GBV-B virus genome could be prepared. It is now known that this obstacle was encountered because the GBV-B genome was believed to be 259 nucleotides shorter than its actual length (Muerhoff *et al.*, 1995; Simons *et al.*, 1995). Others, previous to the inventors, had failed to realize that the 3' sequence of GBV-B was missing from the prior sequences. Without this 3' sequence, it is not possible to prepare an infectious GBV-B molecular clone.

SUMMARY OF THE INVENTION

As discussed above, an infectious molecular clone of GBV-B would be very useful for the development of HCV preventative and therapeutic treatments. The construction of an infectious molecular clone of this virus will require the newly determined 3' sequence to be included in order for the clone to be viable. The inventors have elucidated the previously unrecognized 3' terminal sequence of GBV-B (SEQ ID NO:1). This sequence has been reproducibly recovered from tamarin serum containing GBV-B RNA, in RT-PCR protocols using several different primer sets, and as a fusion with previously reported 5' GBV-B sequences.

The newly identified 3' sequence is not included in published reports of the GBV-B sequence, nor described in patents relating to the original identification of the viral sequence (see U.S. Patent No. 5,807,670 and references therein).

The invention has utility in that the inclusion of the sequence will be necessary for construction of an infectious molecular GBV-B clone. Such clones clearly have the potential to be constructed as chimeras including relevant hepatitis C virus sequences in lieu of the homologous GBV-B sequence, providing unique tools for drug discovery efforts. A full-length molecular clone of GBV-B was constructed, as described in later sections of this specification.

GBV-B can be used as a model for HCV, and the GBV-B genome can be used as the acceptor molecule in the construction of chimeric viral RNAs containing sequences of both HCV and GBV-B. Such studies will allow one to investigate the mechanisms for the different biological properties of these viruses and to discover and investigate potential inhibitors of specific HCV activities (*e.g.*, proteinase) required for HCV replication. However, all this work is dependent upon construction of an infectious clone of GBV-B, which is itself dependent on the incorporation of the correct 3' terminal nucleotide sequence within this clone. GBV-B has unique advantages over HCV in terms of its ability to replicate and cause liver disease in tamarins, which present fewer restrictions to research than chimpanzees, the only nonhuman primate species known to be permissive for HCV.

An infectious molecular clone of GBV-B is expected to have utility in liver-specific gene expression or in gene therapy. This application might be enhanced by the inclusion of HCV genomic sequence in the form of a GBV-B/HCV chimera. Further, an infectious GBV-B/HCV chimera expressing HCV envelope proteins can have utility as a vaccine immunogen for hepatitis C.

A full-length cDNA copy of the GBV-B genome was constructed to contain the newly identified 3' terminal sequences. RNA transcribed from this cDNA copy of the genome would be

infectious when inoculated into the liver of a GBV-B permissive tamarin, giving rise to rescued GBV-B virus particles. A chimeric molecule would then be constructed from this infectious GBV-B clone in which the HCV NS3 proteinase or proteinase/helicase sequence (or other relevant HCV sequences of interest in drug discovery efforts) would be placed in frame in lieu of the homologous GBV-B sequence, and this chimeric cDNA would be used to generate infectious GBV-B/HCV chimeric viruses by intrahepatic inoculation of synthetic RNA in tamarins. Published studies indicate that the GBV-B and HCV proteinases have closely related substrate recognition and cleavage properties, making such chimeras highly likely to be viable. These newly generated chimeric GBV-B/HCV viruses could be used in preclinical testing of candidate HCV NS3 proteinase inhibitors.

Therefore, the present invention encompasses an isolated polynucleotide encoding a 3' sequence of the GBV-B genome. The polynucleotide may include the sequence identified as SEQ ID NO:1. It is contemplated that the polynucleotide may be a DNA molecule or it can be an RNA molecule. It is further contemplated that expression constructs may contain a polynucleotide that has a stretch of contiguous nucleotides from SEQ ID NO:1 and/or SEQ ID NO:2, for example, lengths of 50, 100, 150, 250, 500, 1000, 5000, as well as the entire length of SEQ ID NO:1 or 2, are considered appropriate. Such polynucleotides may also be contained in other constructs of the invention or be used in the methods of the invention. Polynucleotides employing sequences from SEQ ID NO:1 may alternatively contain sequences from SEQ ID NO:2 in the constructs and methods of the present invention. Specific mutations discussed in the Examples and/or figures are included as embodiments of the invention.

The invention is also understood as covering a viral expression construct that includes a polynucleotide encoding a 3' sequence of the GBV-B genome. This expression construct is further understood to contain the sequence identified as SEQ ID NO:1. The present invention contemplates the expression construct as a plasmid or as a virus. Furthermore, the expression construct can express GBV-B sequences; alternatively it may express sequences from a chimeric GBV-B/HCV virus.

The identification and isolation of a 3' sequence of GBV-B additionally provides a method of producing a virus, particularly a full-length virus, by introducing into a host cell a viral expression construct containing a polynucleotide encoding a 3' sequence of GBV-B and by culturing the host cell under conditions permitting production of a virus from the construct. This method can be practiced using a prokaryotic cell as a host cell, or by using a eukaryotic cell as a host cell. Furthermore, the eukaryotic cell can be located within an animal.

A method of producing virus according to the claimed invention can also be employed using a polynucleotide that contains synthetic RNA and/or synthetic DNA. Moreover, a step can be added to the method by also isolating any virus produced from the host cell. The virus can then be purified to homogeneity.

5 In further embodiments, the present invention encompasses an oligonucleotide between about 10 and about 259 consecutive bases of SEQ ID NO:1. This oligonucleotide is contemplated to be about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 50 bases in length, about 100 bases in length, about 150 bases in length, about 200 bases in length, or about 259 bases in length.

10 Additional examples of the claimed invention include a method for identifying a compound active against a viral infection by providing a virus expressed from a viral construct containing a 3' sequence of a GBV-B virus, by contacting the virus with a candidate substance; and by comparing the infectious ability of the virus in the presence of the candidate substance with the infectious ability of the virus in a similar system in the absence of the candidate
15 substance. It is contemplated that the invention can be practiced using GBV-B virus or a GBV-B/HCV chimera. Infectious ability of the virus is comparable to infectivity of the virus. Infectivity can be evaluated by evaluating or assessing the ability of a host cell to become infected by the virus or by the ability of the virus to replicate in the cell. However, it may also be evaluated by virus production, virus infectivity, viral load, or phenotype of the cell such as signs
20 of CPE.

The present invention can also be understood to provide a compound active against a viral infection identified by providing a virus expressed from a viral construct containing a 3' sequence of a GBV-B virus; contacting the virus with a candidate substance; and comparing the infectious ability of the virus in the presence of the candidate substance with the infectious ability
25 of the virus in a similar system in the absence of the candidate substance. In some embodiments an active compound is identified using a GBV-B virus, while in other embodiments an active compound is identified using a GBV-B/HCV chimera. Compounds considered active against viral infection would include, but are not limited to, those compounds that inhibit viral infection or that expedite clearance of the virus.

30 In various embodiments of the invention, a GBV-B polynucleotide may encode a GBV-B/HCV chimera that includes at least part of a 5' NTR sequence derived from a HCV 5' NTR. The 5' NTR may comprise at least one domain, *i.e.*, domain I, II, III and/or IV derived from the 5' NTR of HCV. However, it is specifically contemplated that not all four domains are from

HCV in some embodiments. In certain embodiments, the GVB-B/HCV chimera may include at least domain III of the 5' NTR derived from the 5' NTR of HCV. In yet other embodiments the infectious GBV-B clone may comprise domain III of the 5' NTR of HCV, which may or may not include one or more structural or non-structural genes of HCV also incorporated into the chimeric virus. The portions of the 5' NTR of the GVB-B/HCV chimeras will generally be replaced by analogous sequences from the 5' NTR of HCV. It will be understood that the portions or parts of the 5' NTR of GBV-B that may be replaced include all or part of domain I (including sub-region Ia and Ib of GBV-B), domain II, domain III, domain IV, or any combination thereof. Any combination of 5' NTR domains of GBV-B may be replaced with an analogous region of HCV. In certain embodiments, the replacement of a GBV-B region may be accompanied by the deletion of the 5' NTR GBV-B domain Ib region. In addition, any one, two, or three of the 5' NTR domains of GBV-B may be replaced in any combination with analogous sequences from HCV.

In further embodiments of the invention, a polynucleotide encoding a GBV-B/HCV chimera including a 5' NTR domain III sequence derived from a HCV 5' NTR may be propagated *in vivo*, in particular, in the liver of an appropriate host.

Various other embodiments may include isolated polynucleotides comprising a chimeric GBV-B genome, wherein at least part, but not all of a 5' NTR sequence is derived from a HCV 5' NTR. The polynucleotides may be synthetic RNA, RNA, DNA or the like.

Some embodiments include one or more virus, one or more hepatotropic virus, and/or one or more viral expression constructs comprising a chimeric GBV-B polynucleotides including at least a part of the 5' NTR sequence is derived from a HCV 5' NTR.

Methods of producing a chimeric GBV-B virus encoding at least part of a 5' NTR sequence derived from a HCV 5' NTR sequence comprising introducing into a host cell a viral expression construct comprising a chimeric GBV-B polynucleotide encoding at least part of a 5' NTR sequence derived from a HCV 5' NTR sequence and culturing said host cell under conditions permitting production of a virus from said construct are contemplated. The method may use a host cell that is a eukaryotic cell and the host cell may be in an animal. The method may further include the step of isolating virus from said host cell and in particular purify the virus to homogeneity.

In addition, methods for identifying a compound active against a viral infection comprising are contemplated. The methods may include providing a virus expressed from a viral construct comprising at least part of a 5' NTR derived from a HCV 5' NTR, as described herein;

contacting said virus with a candidate substance; and comparing the infectious ability of the virus in the presence of said candidate substance with the infectious ability of the virus in a similar system in the absence of said candidate substance. Each of the embodiments may use or include any of the 5' NTR chimeras described herein.

5 Other embodiments of the invention may include a compound active against a viral infection identified according to the method described above.

Furthermore, additional methods include providing or administering to a subject or patient in need of a compound active against viral infection the compound active against a viral infection, particularly a flavivirus such as HCV infection.. The patient may have been diagnosed
10 with HCV infection or be at risk for HCV infection, or another flavivirus infection. Alternatively, a subject or patient may be administered a vaccine that immunizes the subject or patient with respect to a flavivirus infection, such as HCV. The invention includes such methods.

It is specifically contemplated that any limitation discussed with respect to one embodiment of the invention may apply to any other embodiment of the invention. Furthermore,
15 any composition of the invention may be used in any method of the invention, and any method of the invention may be used to produce or to utilize any composition of the invention.

The use of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or the alternative are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or."

20 Throughout this application, the term "about" is used to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

As used herein the specification, "a" or "an" may mean one or more, unless clearly indicated otherwise. As used herein in the claim(s), when used in conjunction with the word "comprising," the words "a" or "an" may mean one or more than one. As used herein "another"
25 may mean at least a second or more.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit
30 and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1. Predicted secondary RNA structure at the 3' end of the novel 3' GBV-B sequence. The structure shown is that predicted for the 3' terminal 108 nucleotides of the GBV-B genomic RNA by the MFOLD 3.0 computer program of Zucker and Turner, and has an initial free energy of -48 kcal/mole. The predicted structure contains 3 stem-loops, numbered from the 3' end of the genome and labeled SL-1, SL-2, and SL-3. The structure of SL-1 is highly probable, given its terminal position within the genome. Alternative foldings of SL-2 and SL-3 are possible.

FIG. 2. Illustration of the organization of exemplary chimeric cDNAs containing HCV 5' nontranslated (5' NTR) RNA sequences within the background of a modified (MluI restriction-site containing) GBV-B genetic background.

FIG. 3A and 3B. *In vitro* translation of RLuc reporter transcripts (shown schematically in FIG. 3A) in which the initiation of translation of RLuc is dependent upon the upstream viral or chimeric IRES. FIG. 3B, top panel, shows the SDS-PAGE gel on which the products of translation were separated. FIG. 3B, bottom panel, are the results of the PhosphorImager analysis. Mock = no RNA transcript in the translation mix; and m = size markers.

FIG. 4. Shows the translational activity of RLuc reporter transcripts (normalized to FLuc control transcript) in examples of transfected primary tamarin hepatocytes.

FIG. 5. Shows the replication of an exemplary III^{HC} chimeric RNA following intrahepatic inoculation of synthetic RNA in a GBV-B naïve tamarin (*S. mystax*).

FIG. 6. Shows the replication of an exemplary III^{HC} chimeric virus following intravenous inoculation of a GBV-B naïve tamarin (*S. mystax*) with virus present in the serum of T16444 14 weeks after it had been inoculated with synthetic RNA.

FIG. 7. Shows the sequence differences between chimeric viral RNAs recovered from animals T16444 and T16451, and the parental synthetic III^{HC} RNA.

FIG. 8 Replication of the GB/III^{HC} chimeric virus in primary cultures of tamarin hepatocytes. Viral titers were determined both in infected-cell extracts ("cytosol") and in culture supernatants ("secreted") by a quantitative real-time RT-PCR assay.

FIG. 9 . Replication of the GB/III^{HC} chimeric RNA following intrahepatic inoculation of RNA in a GBV-B naïve tamarin. GBV-B viremia was determined in a quantitative real-time TaqMan® RT-PCR assay targeting GBV-B NS5A sequence (see text for details). Serum ALT activities were monitored on a biweekly basis. Antibodies to NS3 were assayed by an ELISA using recombinant GBV-B NS3 as antigen.

FIG. 10 Mutations identified in the virus isolated at week 14 p.i. in an animal (T16444) inoculated with the GB/III^{HC} chimeric RNA.

FIG. 11 Replication of the GB/III^{HC} chimeric virus following intravenous inoculation in a naïve tamarin.

FIG. 12 Schematic representation of the RNA transcripts of mutated cDNAs derived from the parental GB/III^{HC} construct and their *in vivo* replication capacity following intrahepatic inoculation into GBV-B naïve tamarins.

FIG. 13A and 13B. Replication of the GB/III^{HC}-m3 and GB/III^{HC}-m4 chimeric RNAs following intrahepatic inoculation of RNA in individual naïve tamarins. See Fig. 2 for details relative to monitoring of the infection.

FIG. 14A and 14B Predicted models of the secondary structures of the 59NTRs of GBV-B (FIG. 14A) and HCV (FIG. 14B). (FIG. 14A) Structure proposed for the GBV-B 59NTR by Honda *et al.* (1996 and 1999), each of which is incorporated herein by reference. Major predicted structural domains are labeled I to IV, while individual stem-loops are labeled Ia and are analogous to similarly labeled structures in the HCV 59NTR. Base-pair interactions involving the loop sequence of stem-loop III_f that are predicted to result in a putative RNA pseudoknot are drawn as solid lines. Lightly shaded boxes represent AUG triplets located within the 59NTR, while the solid black box represents the polyprotein translation initiation site. The open box indicates a helical segment at the base of domain II, and open circles represent individual nucleotides that were subjected to site-directed mutagenesis in the studies described here. Unpaired bases within domain II that are conserved in domain II of the HCV structure are shown in boldface type. The 59 and 39 limits of the GBV-B IRES, as determined in this study, are indicated by the arrows. (FIG. 14B) Structure proposed for the HCV 59NTR (Deinhardt *et al.*, 1967, Honda *et al.*, 1996 and 1999, each of which is incorporated herein by reference). There are numerous similarities with the GBV-B structure but no predicted stem-loop structures analogous to the Ib, IIb, and IIc stem-loops of GBV-B.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS**A. GVB-B Virus**

The GBV-B genome structure is very similar to hepatitis C and these viruses share approximately 25% nucleotide identity (Simons *et al.*, 1995; Muerhoff *et al.*, 1995). As indicated above, this makes GBV-B more closely related to HCV than any other known virus. GBV-B genomic RNA is about 9.5 kb in length (Muerhoff *et al.*, 1995) with a structured 5' noncoding region that contains an IRES that shares many structural features with the HCV IRES (Honda *et al.*, 1996; Rijnbrand *et al.*, 1999, each of which is incorporated herein by reference). As in HCV, this IRES drives the cap-independent translation of a long open reading frame. The polyprotein expressed from this reading frame appears to be organized identically to that of HCV, and processed to generate proteins with functions similar to those of HCV (Muerhoff *et al.*, 1995). In fact, the major serine proteinases of these viruses (NS3) have been shown to have similar cleavage specificities (Scarselli *et al.*, 1997). Finally, like HCV and distinct from the pestiviruses, the genomic RNA of GBV-B has a poly(U) tract located near its 3' terminus (Simons *et al.*, 1995; Muerhoff *et al.*, 1995). In addition, unreported sequences located at the extreme 3' end of the genome have been identified. This work indicates that the GBV-B RNA, like that of HCV (HCV (Tanaka *et al.*, 1995; Kolykhalov *et al.*, 1996), terminates in a lengthy run of heterogeneous bases (310 nts in GBV-B) possessing a readily apparent secondary structure

B. Nucleic Acids

The present invention provides a nucleic acid sequence encoding a 3' sequence of the GBV-B genome (SEQ ID NO:1).

It should be clear that the present invention is not limited to the specific nucleic acids disclosed herein. As discussed below, a "3' sequence of the GBV-B genome" may contain a variety of different bases and yet still be functionally indistinguishable from the sequences disclosed herein. Such functionally indistinguishable sequences are likely to maintain the basic structure depicted in FIGs. 1 and 2, which may be used to guide the prediction of viable nucleotide substitutions.

1. Polynucleotides Encoding the 3' Sequence of the GBV-B Genome

A 3' sequence of the GBV-B genome disclosed in SEQ ID NO:1 is one aspect of the present invention. Nucleic acids according to the present invention may encode the 3' sequence of the GBV-B genome set forth in SEQ ID NO:1, the entire GBV-B genome, or any other fragment

of a 3' sequence of the GBV-B genome set forth herein. The nucleic acid may be derived from genomic RNA as cDNA, *i.e.*, cloned directly from the genome of GBV-B. cDNA may also be assembled from synthetic oligonucleotide segments.

It also is contemplated that a 3' sequence of the GBV-B genome may be represented by natural variants that have slightly different nucleic acid sequences but, nonetheless, maintain the same general structure (see FIGs. 1 and 2) and perform the same function in RNA replication.

As used in this application, the term "a nucleic acid encoding a 3' sequence of the GBV-B genome" refers to a nucleic acid molecule that may be isolated free of total viral nucleic acid. In preferred embodiments, the invention concerns nucleic acid sequences essentially as set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12. The term "as set forth in SEQ ID NO:1" means that the nucleic acid sequence substantially corresponds to a portion of SEQ ID NO:1. It is contemplated that the techniques and methods described in this disclosure may apply to any of the sequences contained herein, including SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12.

Allowing for the degeneracy of the genetic code, sequences that have at least about 50%, usually at least about 60%, more usually about 70%, most usually about 80%, preferably at least about 90% and most preferably about 95% of nucleotides that are identical to the nucleotides of SEQ ID NO:1 will be sequences that are "as set forth in SEQ ID NO:1." Sequences that are essentially the same as those set forth in SEQ ID NO:1 may also be functionally defined as sequences that are capable of hybridizing to a nucleic acid segment containing the complement of SEQ ID NO:1 under standard conditions.

The nucleic acid segments and polynucleotides of the present invention include those encoding biologically functional equivalent 3' sequences of the GBV-B genome. Changes designed by man may be introduced through the application of site-directed mutagenesis techniques or may be introduced randomly and screened later for the desired function, as described below.

3' sequence of the GBV-B genome sequences also are provided. Each of the foregoing is included within all aspects of the following description. The present invention concerns cDNA segments reverse transcribed from GBV-B genomic RNA (referred to as "DNA"). As used herein, the term "polynucleotide" refers to an RNA or DNA molecule that may be isolated free of other RNA or DNA of a particular species.

“Isolated substantially away from other coding sequences” means that the 3' sequence of the GBV-B genome forms the significant part of the RNA or DNA segment and that the segment does not contain large portions of naturally-occurring coding RNA or DNA, such as large fragments or other functional genes or cDNA noncoding regions. Of course, this refers to the polynucleotide as originally isolated, and does not exclude genes or coding regions later added to the it by the hand of man.

In certain other embodiments, the invention concerns isolated DNA segments (cDNA segments reverse transcribed from GVB-B genomic RNA) and recombinant vectors that include within their sequence a nucleic acid sequence essentially as set forth in SEQ ID NO:1. The term “essentially as set forth in SEQ ID NO:1” is used in the same sense as described above.

It also will be understood that nucleic acid sequences may include additional residues, such as additional 5' or 3' sequences, and still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological activity. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include additional various non-coding sequences flanking either of the 5' or 3' portions of the coding region, which are known to occur within viral genomes.

Sequences that are essentially the same as those set forth in SEQ ID NO:1 also may be functionally defined as sequences that are capable of hybridizing to a nucleic acid segment containing the complement of SEQ ID NO:1 under relatively stringent conditions. Suitable relatively stringent hybridization conditions will be well known to those of skill in the art.

The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other RNA or DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

For example, nucleic acid fragments may be prepared that include a short contiguous stretch identical to or complementary to SEQ ID NO:1, such as about 15-24 or about 25-34 nucleotides and that are up to about 259 nucleotides being preferred in certain cases. Other stretches of contiguous sequence that may be identical or complementary to any of the sequences disclosed herein, including the SEQ ID NOS. include the following ranges of nucleotides: 50-

9,399, 100-9,000, 150-8,000, 200-7,000, 250-6,000, 300-5,000, 350-4,000, 400-3,000, 450-2,000, 500-1000. RNA and DNA segments with total lengths of about 1,000, about 500, about 200, about 100 and about 50 base pairs in length (including all intermediate lengths) are also contemplated to be useful.

5 In a non-limiting example, one or more nucleic acid constructs may be prepared that include a contiguous stretch of nucleotides identical to or complementary to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12. Such a stretch of nucleotides, or a nucleic acid construct, may be about, or at least about, 3, about 4, about 5, about 10 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39 about 40, about 45, about 50, about 55, about 60, about 65, about 70, about 75, about 80, about 85, about 90, about 95, about 100, about 105, about 15 110, about 115, about 120, about 125, about 130, about 135, about 140, about 145, about 150, about 155, about 160, about 165, about 170, about 175, about 180, about 185, about 190, about 195, about 200, about 210, about 220, about 230, about 240, about 250, about 260, about 270, about 280, about 290, about 300, about 310, about 320, about 330, about 340, about 350, about 360, about 370, about 380, about 390, about 400, about 410, about 420, about 430, about 440, 20 about 450, about 460, about 470, about 480, about 490, about 500, about 510, about 520, about 530, about 540, about 550, about 560, about 570, about 580, about 590, about 600, about 610, about 618, about 650, about 700, about 750, about 1,000, about 2,000, about 3,000, about 4,000, about 5,000, about 6,000, about 7,000, about 8,000, about 9,000, about 9,100, about 9,200, about 9,300, about 9,399, about 9,400, about 9,500, about 9,600, about 9,700, about 9,800, about 25 9,900, about 10,000, about 15,000, about 20,000, about 30,000, about 50,000, about 100,000, about 250,000, about 500,000, about 750,000, to about 1,000,000 nucleotides in length, as well as constructs of greater size, up to and including chromosomal sizes (including all intermediate lengths and intermediate ranges), given the advent of nucleic acids constructs such as a yeast artificial chromosome are known to those of ordinary skill in the art.

30 It will be readily understood that "intermediate lengths," in these contexts means any length between the quoted ranges, such as 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, etc.; 100, 101, 102, 103, etc.; 150, 151, 152, 153, etc.; including all integers through the 200-500;

500-1,000; 1,000-2,000; ranges, up to and including sequences of about 1,001, 1,250, 1,500, and the like.

In certain embodiments, GBV-B polynucleotides include chimeric GBV-B polynucleotides. Chimeric GBV-B polynucleotides may include GBV-B/HCV chimeras. GBV-B/HCV chimeras include, but are not limited to GBV-B polynucleotides in which portions of the GBV-B virus have been replaced with or mutated to resemble analogous sequences from the HCV virus. A GBV-B chimera may comprise all or part of the 5'NTR region of a HCV virus. The polynucleotide may comprise domain I, II, III and/or IV of the 5' NTR derived from a HCV 5'NTR. The polynucleotide may have the 5' NTR domain Ib of GBV-B is deleted. The polynucleotide may comprise domain II and domain III of the 5' NTR derived from a HCV 5'NTR. The polynucleotide may comprise domain II and domain IV of the 5' NTR derived from a HCV 5'NTR. The polynucleotide may comprise domain III and domain IV of the 5' NTR derived from a HCV 5'NTR. The polynucleotide may comprise domain II, domain III and domain IV of the 5' NTR derived from a HCV 5'NTR. Domain Ib of GBV-B may or may not be deleted from the polynucleotide. The polynucleotide may be DNA or RNA. The polynucleotide may further comprising at least part of a structural and/or nonstructural protein coding region of derived from HCV. It will be understood that the term "derived" indicates the origin of the sequence. Thus, the recited domains can be obtained from HCV.

In other embodiments a HCV/GBV-B chimera is contemplated. A HCV/GBV-B chimera will comprise a virus where one or more portions of the HCV virus have been replaced with or mutated to resemble the analogous GBV-B portions of the GBV-B virus.

The various probes and primers designed around the disclosed nucleotide sequences of the present invention may be of any length. By assigning numeric values to a sequence, for example, the first residue is 1, the second residue is 2, *etc.*, an algorithm defining all primers can be proposed:

$$n \text{ to } n + y$$

where n is an integer from 1 to the last number of the sequence and y is the length of the primer minus one, where $n + y$ does not exceed the last number of the sequence. Thus, for a 20-mer, the probes correspond to bases 1 to 20, 2 to 21, 3 to 22 ... and so on. For a 30-mer, the probes correspond to bases 1 to 30, 2 to 31, 3 to 32 ... and so on. For a 35-mer, the probes correspond to bases 1 to 35, 2 to 36, 3 to 37 ... and so on.

2. Oligonucleotide Probes and Primers

Naturally, the present invention also encompasses RNA and DNA segments that are complementary, or essentially complementary, to the sequence set forth in SEQ ID NO:1. Nucleic

acid sequences that are "complementary" are those that are capable of base-pairing according to the standard Watson-Crick complementary rules. As used herein, the term "complementary sequences" means nucleic acid sequences that are substantially complementary, as may be assessed by the same nucleotide comparison set forth above, or as defined as being capable of hybridizing to the nucleic acid segment of SEQ ID NO. 1 under relatively stringent conditions such as those described herein. Such sequences may encode the entire 3' sequence of the GBV-B genome or functional or non-functional fragments thereof.

Alternatively, the hybridizing segments may be shorter oligonucleotides. Sequences of 17 bases long should occur only once in the human genome and, therefore, suffice to specify a unique target sequence. Although shorter oligomers are easier to make and increase *in vivo* accessibility, numerous other factors are involved in determining the specificity of hybridization. Both binding affinity and sequence specificity of an oligonucleotide to its complementary target increases with increasing length. It is contemplated that exemplary oligonucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225 or more base pairs will be used, although others are contemplated. Longer polynucleotides encoding 250, 500, 1000, 1212, 1500, 2000, 2500, 3000 or 3431 bases and longer are contemplated as well. Such oligonucleotides will find use, for example, as probes in Southern and Northern blots and as primers in amplification reactions.

Suitable hybridization conditions will be well known to those of skill in the art. In certain applications, for example, substitution of amino acids by site-directed mutagenesis, it is appreciated that lower stringency conditions are required. Under these conditions, hybridization may occur even though the sequences of probe and target strand are not perfectly complementary but are mismatched at one or more positions. Conditions may be rendered less stringent by increasing salt concentration and decreasing temperature. For example, a medium stringency condition could be provided by about 0.1 to 0.25 M NaCl at temperatures of about 37°C to about 55°C, while a low stringency condition could be provided by about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20°C to about 55°C. Thus, hybridization conditions can be readily manipulated and thus will generally be a method of choice depending on the desired results.

In other embodiments, hybridization may be achieved under conditions of, for example, 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl₂, 10 mM dithiothreitol, at temperatures between approximately 20°C to about 37°C. Other hybridization conditions utilized could include approximately 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, at temperatures ranging

from approximately 40°C to about 72°C. Formamide and SDS also may be used to alter the hybridization conditions.

One method of using probes and primers of the present invention is in the search for other viral sequences related to GBV-B or, more particularly, homologs of the GBV-B sequence. By varying the stringency of hybridization, and the region of the probe, different degrees of homology may be discovered.

Another way of exploiting probes and primers of the present invention is in site-directed, or site-specific, mutagenesis. The technique provides a ready ability to prepare and test sequence variants, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into complementary DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences that encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered.

The technique typically employs a bacteriophage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage vectors are commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis, which eliminates the step of transferring the gene of interest from a phage to a plasmid.

In general, site-directed mutagenesis is performed by first obtaining a single-stranded vector, or melting of two strands of a double stranded vector which includes within its sequence a DNA sequence encoding the desired protein. An oligonucleotide primer bearing the desired mutated sequence is synthetically prepared. This primer is then annealed with the single-stranded DNA preparation, taking into account the degree of mismatch when selecting hybridization conditions, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected that include recombinant vectors bearing the mutated sequence arrangement. There are newer and simpler site-directed mutagenesis

techniques that can also be employed for this purpose. These include procedures marketed in kit form that are readily available to one of ordinary skill in the art.

The preparation of sequence variants of the selected gene using site-directed mutagenesis is provided as a means of producing potentially useful species and is not meant to be limiting, as there are other ways in which sequence variants of genes may be obtained. For example, recombinant vectors encoding the desired gene may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

3. Antisense Constructs

In certain embodiments of the invention, the use of antisense constructs of the 3' sequence of the GBV-B genome is contemplated.

Antisense methodology takes advantage of the fact that nucleic acids tend to pair with "complementary" sequences. By complementary, it is meant that polynucleotides are those which are capable of base-pairing according to the standard Watson-Crick complementarity rules. That is, the larger purines will base pair with the smaller pyrimidines to form combinations of guanine paired with cytosine (G:C) and adenine paired with either thymine (A:T) in the case of DNA, or adenine paired with uracil (A:U) in the case of RNA. Inclusion of less common bases such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others in hybridizing sequences does not interfere with pairing.

Targeting double-stranded (ds) DNA with polynucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation. Antisense polynucleotides, when introduced into a target cell, specifically bind to their target polynucleotide and interfere with transcription, RNA processing, transport, translation and/or stability. Antisense RNA constructs, or DNA encoding such antisense RNAs, may be employed to inhibit gene transcription or translation or both within a host cell, either *in vitro* or *in vivo*, such as within a host animal, including a human subject.

Antisense constructs could be used to block early steps in the replication of GBV-B and related viruses, by annealing to 3' terminal sequences and blocking their role in negative-strand initiation.

As stated above, "complementary" or "antisense" means polynucleotide sequences that are substantially complementary over their entire length and have very few base mismatches. For example, sequences of 15 bases in length may be termed complementary when they have complementary nucleotides at 13 or 14 positions. Naturally, sequences which are completely complementary will be sequences which are entirely complementary throughout their entire

length and have no base mismatches. Other sequences with lower degrees of homology also are contemplated. For example, an antisense construct which has limited regions of high homology but also contains a non-homologous region (*e.g.*, ribozyme; see below) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

4. Amplification and PCR™

The present invention utilizes amplification techniques in a number of its embodiments. Nucleic acids used as a template for amplification are isolated from cells contained in the biological sample, according to standard methodologies (Sambrook *et al.*, 1989). The nucleic acid may be genomic DNA or RNA or fractionated or whole cell RNA. Where RNA is used, it may be desired to convert the RNA to a complementary DNA using reverse transcriptase (RT). In one embodiment, the RNA is genomic RNA and is used directly as the template for amplification. In others, genomic RNA is first converted to a complementary DNA sequence (cDNA) and this product is amplified according to protocols described below.

Pairs of primers that selectively hybridize to nucleic acids corresponding to GBV-B sequences are contacted with the isolated nucleic acid under conditions that permit selective hybridization. The term "primer," as defined herein, is meant to encompass any nucleic acid that is capable of priming the synthesis of a nascent nucleic acid in a template-dependent process. Typically, primers are oligonucleotides from ten to twenty base pairs in length, but longer sequences can be employed. Primers may be provided in double-stranded or single-stranded form, although the single-stranded form is preferred.

Once hybridized, the nucleic acid:primer complex is contacted with one or more enzymes that facilitate template-dependent nucleic acid synthesis. Multiple rounds of amplification, also referred to as "cycles," are conducted until a sufficient amount of amplification product is produced.

Next, the amplification product is detected. In certain applications, the detection may be performed by visual means. Alternatively, the detection may involve indirect identification of the product via chemiluminescence, radioactive scintigraphy of incorporated radiolabel or fluorescent label or even via a system using electrical or thermal impulse signals.

A number of template dependent processes are available to amplify the marker sequences present in a given template sample. One of the best known amplification methods is the polymerase chain reaction (referred to as PCR™) which is described in detail in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,800,159, and each incorporated herein by reference in entirety.

Briefly, in PCR™, two or more primer sequences are prepared that are complementary to regions on opposite complementary strands of the marker sequence. An excess of deoxynucleoside triphosphates are added to a reaction mixture along with a DNA polymerase, *e.g.*, *Taq* polymerase. If the marker sequence is present in a sample, the primers will bind to the marker and the polymerase will cause the primers to be extended along the marker sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the marker to form reaction products, excess primers will bind to the marker and to the reaction products and the process is repeated.

A reverse transcriptase PCR™ amplification procedure may be performed in order to quantify the amount of mRNA amplified. Methods of reverse transcribing RNA into cDNA are well known and described in Sambrook *et al.*, 1989. Alternative methods for reverse transcription utilize thermostable, RNA-dependent DNA polymerases. These methods are described in WO 90/07641, filed December 21, 1990, incorporated herein by reference. Polymerase chain reaction methodologies are well known in the art.

Another method for amplification is the ligase chain reaction ("LCR"), disclosed in EPA No. 320 308, incorporated herein by reference in its entirety. U.S. Patent 4,883,750 describes a method similar to LCR for binding probe pairs to a target sequence.

Qbeta Replicase, described in PCT Application No. PCT/US87/00880, incorporated herein by reference, also may be used as still another amplification method in the present invention.

An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[alpha-thio]-triphosphates in one strand of a restriction site also may be useful in the amplification of nucleic acids in the present invention.

Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, *i.e.*, nick translation. A similar method, called Repair Chain Reaction (RCR), involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA. Target specific sequences also can be detected using a cyclic probe reaction (CPR). In CPR, a probe having 3' and 5' sequences of non-specific DNA and a middle sequence of specific RNA is hybridized to DNA that is present in a sample. Upon hybridization, the reaction is treated with RNase H, and

the products of the probe identified as distinctive products that are released after digestion. The original template is annealed to another cycling probe and the reaction is repeated.

5 Still another amplification methods described in GB Application No. 2 202 328, and in PCT Application No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template- and enzyme-dependent synthesis. The primers may be modified by labeling with a capture moiety (*e.g.*, biotin) and/or a detector moiety (*e.g.*, enzyme). In the latter application, an excess of labeled probes are added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, 10 the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS), including nucleic acid sequence based amplification (NASBA) and 3SR Gingeras *et al.*, PCT Application WO 88/10315, incorporated herein by reference.

15 Davey *et al.*, EPA No. 329 822 (incorporated herein by reference in its entirety) disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention.

20 Miller *et al.*, PCT Application WO 89/06700 (incorporated herein by reference in its entirety) disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target ssDNA followed by transcription of many RNA copies of the sequence. This scheme is not cyclic, *i.e.*, new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" and "one-sided PCR" (Frohman, 1990 incorporated by reference).

25 Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide", thereby amplifying the di-oligonucleotide, also may be used in the amplification step of the present invention.

Following any amplification, it may be desirable to separate the amplification product from the template and the excess primer for the purpose of determining whether specific 30 amplification has occurred. In one embodiment, amplification products are separated by agarose, agarose-acrylamide or polyacrylamide gel electrophoresis using standard methods. See Sambrook *et al.*, 1989.

Alternatively, chromatographic techniques may be employed to effect separation. There are many kinds of chromatography that may be used in the present invention: adsorption, partition, ion-exchange and molecular sieve, and many specialized techniques for using them including column, paper, thin-layer and gas chromatography.

5 Amplification products must be visualized in order to confirm amplification of the marker sequences. One typical visualization method involves staining of a gel with ethidium bromide and visualization under UV light. Alternatively, if the amplification products are integrally labeled with radio- or fluorometrically-labeled nucleotides, the amplification products can then be exposed to x-ray film or visualized under the appropriate stimulating spectra,
10 following separation.

In one embodiment, visualization is achieved indirectly. Following separation of amplification products, a labeled nucleic acid probe is brought into contact with the amplified marker sequence. The probe preferably is conjugated to a chromophore but may be radiolabeled. In another embodiment, the probe is conjugated to a binding partner, such as an antibody or
15 biotin, and the other member of the binding pair carries a detectable moiety.

In one embodiment, detection is by Southern blotting and hybridization with a labeled probe. The techniques involved in Southern blotting are well known to those of skill in the art and can be found in many standard books on molecular protocols. See Sambrook *et al.*, 1989. Briefly, amplification products are separated by gel electrophoresis. The gel is then contacted
20 with a membrane, such as nitrocellulose or nylon, permitting transfer of the nucleic acid and non-covalent binding. Subsequently, the membrane is incubated with a chromophore-conjugated probe that is capable of hybridizing with a target amplification product. Detection is by exposure of the membrane to x-ray film or ion-emitting detection devices.

One example of the foregoing is described in U.S. Patent 5,279,721, incorporated by
25 reference herein, which discloses an apparatus and method for the automated electrophoresis and transfer of nucleic acids. The apparatus permits electrophoresis and blotting without external manipulation of the gel and is ideally suited to carrying out methods according to the present invention.

5. Expression Constructs

30 In some embodiments of the present invention, an expression construct that encodes a 3' sequence of GBV-B is utilized. The term "expression construct" is meant to include any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated

into a protein, but it need not be. Expression includes both transcription of a gene and translation of mRNA into a gene product. Expression may also include only transcription of the nucleic acid encoding a gene of interest.

5 In some constructs, the nucleic acid encoding a gene product is under transcriptional control of promoter and/or enhancer. The term promoter will be used here to refer to a group of transcriptional control modules that are clustered around the initiation site for RNA polymerase II. Much of the thinking about how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (tk) and SV40 early transcription units. These studies have shown that promoters are composed of discrete functional modules, 10 each consisting of approximately 7-20 bp of nucleic acids, and containing one or more recognition sites for transcriptional activator or repressor proteins.

At least one module in each promoter functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene 15 and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start 20 site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

25 Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of nucleic acids with enhancer activity are organized much like promoters. That is, they are composed of many individual elements, each of which binds to 30 one or more transcriptional proteins.

The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or

more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

6. Host Cells and Permissive Cells

As used herein, the terms "cell," "cell line," and "cell culture" may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of the present invention, "host cell" refers to a prokaryotic or eukaryotic cell, and it includes any transformable organisms that is capable of replicating a vector or virus and/or expressing viral proteins. A host cell can, and has been, used as a recipient for vectors, including viral vectors. A host cell may be "transfected" or "transformed," which refers to a process by which exogenous nucleic acid is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny. A "permissive cell" refers to a cell that supports the replication of a given virus and consequently undergoes cell lysis. In the context of the present invention, such a virus would include HCV, GBV-B, or other hepatitis viruses. In a "nonpermissive cell," productive infection does not result, but the cell may become stably transformed. In some embodiments, methods employ permissive cells that are a cell line derived from liver cells (liver cell line).

Host cells may be derived from prokaryotes or eukaryotes, depending upon whether the desired result is replication of the vector or expression of part or all of the vector-encoded nucleic acid sequences. Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials (www.atcc.org). An appropriate host can be determined by one of skill in the art based on the vector backbone and the desired result. A plasmid or cosmid, for example, can be introduced into a prokaryote host cell for replication of many vectors. Bacterial cells used as host cells for vector replication and/or expression include DH5 α , JM109, and KC8, as well as a number of commercially available bacterial hosts such as SURE[®] Competent Cells and SOLOPACK[™] Gold Cells (STRATAGENE[®], La Jolla). Alternatively, bacterial cells such as *E. coli* LE392 could be used as host cells for phage viruses.

Examples of eukaryotic host cells for replication and/or expression of a vector include HeLa, NIH3T3, Jurkat, 293, Cos, CHO, Saos, and PC12. Many host cells from various cell types and organisms are available and would be known to one of skill in the art. Similarly, a viral

vector or virus or virus particle may be used in conjunction with either a eukaryotic or prokaryotic host cell, particularly one that is permissive for replication or expression of the vector. It is contemplated that the present invention includes vectors composed of viral sequences, viruses, and viral particles in the methods of the present invention, and that they may be used interchangeably in these methods, depending on their utility.

Some vectors may employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to maintain them and to permit replication of a vector. Also understood and known are techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and their cognate polypeptides, proteins, or peptides.

7. Pharmaceutical Compositions

The present invention encompasses the use of a 3' sequence of GBV-B in the production of or use as a vaccine to combat HCV infection. Compositions of the present invention comprise an effective amount of GBV-B clone as a therapeutic dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. The phrases "pharmaceutically or pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. For human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

The biological material should be extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle, where appropriate. The active compounds will then generally be formulated for parenteral administration, *e.g.*, formulated for injection via the intravenous, intramuscular, sub-cutaneous, intralesional, or even intraperitoneal routes. The preparation of an aqueous composition that contains GVB-B nucleic acid sequences as an active component or ingredient will be known to

those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and the preparations can also be emulsified.

5 The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against
10 the contaminating action of microorganisms, such as bacteria and fungi.

 Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a
15 preservative to prevent the growth of microorganisms.

 A GBV-B clone of the present invention can be formulated into a composition in a neutral or salt form. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric,
20 mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. In terms of using peptide therapeutics as active ingredients, the technology of U.S. Patents 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, each
25 incorporated herein by reference, may be used.

 The carrier also can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size
30 in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged

absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

5 Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active
10 ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The preparation of more, or highly, concentrated solutions for direct injection is also contemplated, where the use of DMSO as solvent is envisioned to result in extremely rapid penetration, delivering high concentrations of the active agents to a small area.

15 Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above, but drug release capsules and the like also can be employed.

20 For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of
25 infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

30 In addition to the compounds formulated for parenteral administration, such as intravenous or intramuscular injection, other pharmaceutically acceptable forms include, *e.g.*, tablets or other solids for oral administration; liposomal formulations; time release capsules; and any other form currently used, including cremes.

Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders. In certain defined embodiments, oral pharmaceutical compositions will comprise an inert diluent or assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 75% of the weight of the unit, or preferably between 25-60%. The amount of active compounds in such therapeutically useful compositions is such that a suitable dosage will be obtained.

The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compounds sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor.

C. Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

EXAMPLE 1

Elucidation of a 3' Sequence of the GBV-B Genome

The inventors elucidated the previously unrecognized 3' terminal sequence of GBV-B (SEQ ID NO 1). This sequence was reproducibly recovered from tamarin serum containing GBV-B RNA, in RT-PCR nucleic acid amplification procedures using several different primer sets, and as a fusion with previously reported 5' GBV-B sequences.

There is information in the published literature reporting the putative sequences of the 5' and 3' termini of the GBV-B genome. The nucleic acid sequences of these termini were reportedly determined by ligating the ends of the viral RNA together, amplifying the sequence in the region of the resulting junction by reverse-transcription polymerase chain reaction (RT-PCR), and sequencing of the cDNA amplification product across the junction. However, the inventors believed these results required confirmation. Of particular concern was the fact that the 3' terminus appeared to be shorter than the equivalent region of other viruses in the family *Flaviviridae* (especially within the genus *Hepacivirus*) and that the reported 3' sequence lacked a defined RNA hairpin structure such as those present in these related viruses. Additional novel sequences at the 3' end of the GBV-B genome were investigated using a serum sample collected from a tamarin that was experimentally infected with virus. Amplification was used to determine the sequence of the 3' end.

First, serum (50 μ L) known to contain GBV-B RNA by RT-PCR assay was extracted with Trizol, and the RNA was washed and dried. A synthetic oligonucleotide was then ligated to the 3' end of the viral RNA. The oligonucleotide, 5'-AATTCGGCCCTGCAGGCCACAACAGTC-3', which was phosphorylated at the 5' end and chemically blocked at the 3' end, was ligated to the RNA essentially using the method described by Kolykhalov *et al.* (Behrens *et al.*, 1996). The RNA was initially dissolved in DMSO and the following additions were made: Tris-Cl, pH 7.5 (10 mM), $MgCl_2$ (10 mM), DTT (5 mM), hexamine cobalt chloride (1 mM), 10 pmol oligo and 8 U T4 ligase. The final concentration of DMSO was 30% in a final volume of 10 μ L. The ligation reaction was incubated for 4 or 20 hours at 19° C. 1 μ L of the ligation reaction was used directly to make cDNA, using a primer complementary to the ligated oligonucleotide and the Superscript 2 system, in a final volume of 15 μ L. 1 μ L of cDNA was amplified using the Advantage cDNA system (Clontech) and two additional oligonucleotide primers. These primers included one that was complementary to the ligated oligonucleotide (*i.e.*, "negative sense") and a positive-sense primer located near the 3' end of the reported GBV-B sequence. A product approximately 290 bases in length was obtained,

and this was gel purified and directly sequenced. Sequencing was done in both directions using the oligonucleotide primers employed for the amplification; 259 bases that had not been previously reported were identified as fused to the sequence that had been previously described as the 3' terminus of the viral genome.

5 To ensure that this novel 3' sequence from viral RNA could be reproducibly amplified, an additional 10 µL of infected tamarin serum was extracted using Trizol. cDNA was prepared by reverse transcription using an oligonucleotide primer complementary to the penultimate 3' 25 bases of the novel sequence. Amplification was then done by PCR using the primer previously utilized for cDNA synthesis and a positive-sense primer mapping within the previously published
10 GBV-B sequence. In the initial studies, although a product was readily detected, DNA sequencing showed that this product was missing all of the sequence distal to the poly-U tract. Carrying out the cDNA synthesis in the presence of DMSO circumvented this problem. A cDNA product of approximately 290 bases was obtained. This was sequenced and shown to consist of the 5' primer, 20 bases of the published GBV-B sequence, and 259 bases of the novel sequence
15 obtained in the preceding studies and containing the sequence of the 3' primer. The sequence of the 3' end of GBV-B is shown in SEQ ID NO: 1 (FIG. 1). The possible secondary RNA structure for this region is shown in FIG. 2, as predicted by a computer-based RNA folding program. The presence of a predicted hairpin structure at the extreme 3' end of this novel sequence is consistent with its location at the 3' terminus of the viral RNA.

20 The GBV-B cDNA (synthesis described above) was used as a template for PCR amplification of the 3' 1553 nucleotides (nts) of the GBV-B genome. This PCR amplification product was gel purified and cloned into plasmid DNA using the "Perfectly Blunt Cloning Kit" (Novagene).

EXAMPLE 2

25 Construction of an Infectious GBV-B Clone

The elucidation of a 3' sequence of the GBV-B genome will allow those of skill in the art to construct and validate an infectious molecular clone of GBV-B. This will be done using the following procedures.

A full-length cDNA copy of the GBV-B genome containing the newly identified 3'
30 terminal sequences was constructed. RNA transcribed from this cDNA copy of the genome will be infectious when inoculated into the liver of a GBV-B permissive tamarin, giving rise to rescued GBV-B virus particles.

A 1:1000 dilution of GBV-B infectious tamarin serum was obtained. This material was used as a source of viral RNA for the amplification of GBV-B nucleic acid sequences by reverse-transcription polymerase chain reaction. For amplification of previously reported segments of the GBV-B genome, 250 μ L of the diluted serum was extracted with Trizol using the manufacturer's instructions. The final RNA pellet was dissolved in 10 μ L of a 100 mM DTT buffer containing 5% RNasin. This material was converted into cDNA using Superscript 2 reverse transcriptase and oligonucleotide primers designed to be complementary to the reported GBV-B RNA sequence and to contain unique restriction sites. This cDNA was amplified using the Advantage cDNA kit (Clontech) employing the cDNA primer (negative sense as the downstream primer and a similar positive-sense upstream primer, again containing a unique restriction site. The published sequence of GBV-B allowed for the selection of primers in convenient areas of the genome containing unique restriction sites. Using this general strategy, the inventors amplified segments of the reported GBV-B genome representing: (1) nucleotides (nts) 1-1988, using an upstream primer containing a T7 RNA polymerase promoter and a BamH1 site upstream of nt 1, and a downstream primer containing a unique EcoR1 site (nt 1978); (2) nts 1968-5337, using a downstream primer containing a unique Cla1 site at position 5027; (3) nts 5317- 7837, using a downstream primer containing a Sal1 site at nt 7847; and, (4) nts 7837-9143, using a downstream primer containing an added Xho1 site. It was found necessary to use different PCR conditions for each primer set.

The RT-PCR products generated in these reactions were cloned into plasmid DNA after gel purification, using the "Perfectly Blunt Cloning Kit" (Novagene). Ten bacterial colonies from each of the four RT-PCR products were analyzed for insert size by restriction endonuclease digestion using EcoR1, the sites for this enzyme being located on either side of the insert in the resulting plasmids. For three of the RT-PCR amplicons, 9 of 10 colonies contained plasmids with the correct size insert. The EcoR1-Cla1 amplicon generated only 1/10 colonies with a correct size insert. Thus, 30 additional colonies were examined, yielding two more clones with insert of the correct size. For each of these plasmids, simple restriction patterns were obtained using two restriction enzymes. As these appeared to be correct, the plasmid DNAs were subjected to sequencing using an ABI automatic sequencer.

EXAMPLE 3

Nucleotide sequence of the cloned GBV-B cDNA

The 5' region of the cloned sequence revealed a relatively long nontranslated region corresponding to the published sequence of the GBV-B 5'NTR, which includes an IRES. This

region was followed by a long open reading frame. Near the 3' end of the genome a poly-U tract was identified; however, this was shorter than the published 3' homopolymeric poly-U region. The sequence from these clones was compared with those in the GenBank database (Accession U22304, "Hepatitis GB virus B polypeptide complete genome"). Twenty-two nucleotide differences were identified, of which 14 gave rise to amino acid changes (Table 1). In order to determine whether these changes were genuine or RT-PCR artifacts, which could have been introduced due to the very small amount of material from which these sequences were amplified, segments of the genome containing these changes were reamplified using a serum sample from an independently infected tamarin. Of the 14 changes noted in the original cDNA clones, 12 were not present in these newly amplified sequences and thus were probably RT-PCR artifacts (Table 2.). A particularly interesting difference from the published GenBank sequence, however, which was present in both the original clones as well as a repeat amplification, was a two-nucleotide substitution that obliterated the Sal I site present in the published sequence.

Table 1. Differences in the amino acid sequences of GBV-B cDNA clones and the GBV-B sequence reported by Simons *et al.* (1995).

GBV-B Protein	AMINO ACID Δ FROM ABBOTT SEQUENCE	RT-PCR Products from Tamarin 12024 (PCR Reaction #)
Core	G ₉₉ →S	G (38.1a)
E1	V ₃₉₅ →I	V (40.2a)
E2	D ₇₀₃ →N	D (42.3a)
E2	P ₇₀₆ →Q	H (42.3a)
E2	A ₇₂₈ →V	A (42.3a)
NS2	L ₇₉₁ →F	F(42.3a)
NS2	T ₈₀₄ →A	A(42.3a)
NS5A	L ₁₉₉₀ →M	L (46.5a)
NS5A	I ₂₀₈₂ →T	I (46.5a)
NS5A	S ₂₁₇₄ →P	(not done)
NS5A	G ₂₂₂₈ →E	E (48.6a)
NS5A	T ₂₂₃₃ →S	T (48.6a)
NS5A	A ₂₂₃₆ →V	A (48.6a)
NS5B	V ₂₈₃₃ →I	V (50.7a)

EXAMPLE 4

Construction of a full-length GBV-B cDNA clone.

The four GBV-B cDNA inserts described above were cloned into Bluescript ks+ using unique restriction sites. Since the unique SalI site that was reported to be present in the published GBV-B sequence (nt position 7847) was absent in these cDNA clones, this restriction site was created by engineering two silent nucleotide changes using the "Quick Change" mutagenesis system (Stratagene). Although the most 5' clones (nucleotides 1-7847) could be readily constructed, attempts to add the remaining 3' clones were unsuccessful due to rearrangements and deletions. This problem was overcome by use of pACNR1180, a plasmid that had been used to construct an infectious clone of yellow fever virus. Finally, the most 3' 771 nucleotides of GBV-B were excised from the plasmid containing the novel, previously unreported 3' sequence, and inserted into the truncated assembled GBV-B cDNA construct to complete the 3' end. The 3' terminus of this full-length cDNA was then subjected to DNA sequencing to confirm its integrity. Extensive restriction digests indicated that this construct had the characteristics of a full-length cDNA copy of GBV-B virus. Because there is not yet an understanding of which cultured cells (if any) might be permissive for GBV-B replication, the infectivity of the synthetic GBV-B RNA will be assessed by injecting the RNA directly into the liver of a susceptible tamarin.

Alternatively, an infectious full-length clone can be produced by the following protocol. A plasmid will be made containing a cassette including the 5' and 3' ends of the virus flanked by appropriate restriction sites. These constructs have been shown to efficiently translate reporter genes, with transcription taking place via a T7 promoter placed immediately upstream of the 5'NTR (*e.g.*, see Rijnbrand *et al.*, 1999). The major portion of the GBV-B genome would then be amplified by long range RT-PCR. This method is now well established for hepatitis C virus and other flaviviruses (Teller *et al.*, 1996), and it has been used successfully also to amplify rhinovirus RNA. Briefly this technique uses "Superscript" reverse transcriptase to synthesize cDNA and a mixture of "KlenTaq 1", and "DeepVent" polymerases to amplify this cDNA. Primers that can be used will contain restriction sites to allow cloning of the RT-PCR products into the cassette vector. After being transformed into suitable competent bacteria, extensive restriction analysis will enable us to determine which clones contain inserts that are of full length and which have a high probability of being correct. Apparent full-length clones will be analyzed further by coupled transcription-translation using the Promega "TnT" system, with the addition of microsomal membranes to allow the cleavage of the structural proteins by cellular signalase

enzymes. Clones which appear correct by restriction analysis and which produce GBV-B proteins, in particular the protein coded for by the extreme 3' end of the genome, NS5B, will be selected, and RNA will be transcribed from these clones using the Ambion MegaScript system. "Correct" looking clones (>10) can be injected directly into a tamarin liver at several sites. A successful infection will be determined as described below. If a positive signal is detected the entire genome will be amplified and sequenced to determine which plasmid the virus originated from.

EXAMPLE 5

Rescue of Infectious GVB-B

Infectious GBV-B will be rescued from synthetic genome-length RNA following its injection into the liver of tamarins (*Saguinus sp.*). In past studies, HAV from synthetic RNA in owl monkeys has been recovered (*Aotus trivirgatus*) (Shaffer *et al.*, 1995), and more recently, the recovery of virus from a chimpanzee injected intrahepatically with RNA transcribed from a full-length genotype 1b HCV cDNA clone was reported (Beard *et al.*, 1999).

RNA will be prepared for these studies using the T7 MegaScript kit (Ambion) and a total of 10 µg of plasmid DNA as template. An aliquot of the reaction products will be utilized to ensure the integrity of the RNA by electrophoresis in agarose-formaldehyde gels. The remainder of the transcription reaction mix will be frozen at -80C° until its injection, without further purification, into the liver of a tamarin. Because of the small size of the tamarin, the RNA will be injected under direct visualization following a limited incision and exposure of the liver. Under similar conditions, in other primate species, RT-PCR-detectable viral RNA or cDNA has not been detected in serum samples collected within days of this procedure in the absence of viral replication (Kolykhalov *et al.*, 1997; Yanagi *et al.*, 1998; Beard *et al.*, 1999). Thus, the appearance of RNA in serum collected subsequently from these tamarins will be strong evidence for the replication competence of the synthetic RNA. Serum will be collected weekly for six weeks, then every other week for an additional 6 weeks from inoculated animals. In addition to RT-PCR for detection of viral RNA (see FIG. 2B), alanine aminotransferase (ALT) levels will be measured as an indicator of liver injury and to assess liver histology in punch biopsies taken at the time of ALT elevation. Maximum viremia and an acute phase ALT response is expected to occur around 14-28 days post-inoculation of infectious RNA (Simons *et al.*, 1995; Schlauder *et al.*, 1995; Karayiannis *et al.*, 1989). Transfections will be considered to have failed to give rise to infectious virus if RNA is not detected in the serum within 12 weeks of inoculation.

Successfully infected animals will be followed with twice weekly bleeds until resolution of the viremia, or for 6 months, whichever is longest.

EXAMPLE 6

Construction of GBV-B/HCV Chimeras

The GBV-B genome can be used as the acceptor molecule in the construction of chimeric viral RNAs containing sequences of both HCV and GBV-B. Such constructs will allow one to investigate the mechanisms for the different biological properties of these viruses and to discover and investigate potential inhibitors of specific HCV activities (*e.g.*, proteinase) required for HCV replication. Different classes of chimeric viruses are contemplated. These include: (a) replacement of the GBV-B IRES with that of HCV; and (b) replacement of the NS3 major serine proteinase and helicase, and (c) the replacement of the NS5B RNA-dependent RNA polymerase with the homologous proteins of HCV.

The chimeric constructs described in the following sections will be made by PCR mutagenesis, using high fidelity polymerases and oligonucleotide primers designed to include the specific fusions of GBV-B and HCV sequences (Landt *et al.*, 1990). First round PCR reactions will create the desired fusion, and generate a new "primer" to be used in a second PCR reaction spanning the region to a convenient unique restriction site. PCR cycles will be kept to the minimum number necessary for successful amplification, and all segments of viral sequence that are amplified by PCR will be subjected to DNA sequencing to exclude the presence of unwanted PCR-introduced errors. Sequencing will be accomplished at UTMB's core Recombinant DNA Laboratory. Amplified segments will be kept to the minimum by the exchange of cloned cDNA segments spanning convenient restriction sites in subgenomic clones, and where necessary PCR artifacts can be corrected by site-directed mutagenesis (QuickChange mutagenesis kit, Stratagene).

A number of viable positive-strand RNA virus chimeras have been constructed previously in which IRES elements have been swapped between different viruses. Most of these chimeras have involved the exchange of IRES elements between picornaviruses. Others have been successful in constructing viable poliovirus chimeras containing the HCV IRES in place of the native poliovirus IRES (Zhao *et al.*, 1999; Lu and Wimmer, 1996). A similar rhinovirus 14 chimera containing the HCV IRES has been constructed, although its replication phenotype is not as robust as the poliovirus chimera described by Lu and Wimmer (Lu and Wimmer, 1996). More importantly, Frolov *et al.* (Frolov *et al.*, 1998) recently reported chimeric flaviviruses in

which the HCV IRES was inserted into the genetic background of a pestivirus, bovine viral diarrhea virus (BVDV) in lieu of the homologous BVDV sequence. Although these viable chimeric polioviruses and pestiviruses replicate in cell cultures, they are poor surrogates for HCV in animal models as neither virus is hepatotropic or causes liver disease. Importantly, Frolov *et al.* (Frolov *et al.*, 1998) demonstrated quite convincingly that the requirement for cis-acting replication signals at the 5' terminus of the pestivirus genome was limited to a short tetranucleotide sequence. This requirement presumably reflects the need for the complement of this sequence at the 3' end of the negative strand during initiation of positive-strand RNA synthesis. The work of Frolov *et al.* shows that the IRES of BVDV does not contain necessary replication signals, or that if these are present within the BVDV IRES they can be complemented with similar signals in either the HCV or encephalomyocarditis virus (EMCV, a picornavirus) IRES sequence. Since GBV-B and HCV are more closely related to each other than BVDV and HCV, these studies provide strong support for the viability of chimeras containing the HCV IRES in the background of GBV-B.

Construction of a viable IRES chimera will be enhanced by studies that have documented the sequence requirements and secondary structures of the IRES elements of both HCV and GBV-B (*see* Lemon and Honda, 1997; Honda *et al.*, 1996; Rijnbrand *et al.*, 1999). To a considerable extent, the work of Frolov *et al.* (Frolov *et al.*, 1998) was guided by studies of the HCV IRES structure. More recently, these studies have been extended to include a detailed mutational analysis of the GBV-B IRES. The results of these studies indicate that the functional IRES of GBV-B extends from the 5' end of structural domain II (nt 62) to the initiator AUG codon (nt 446). This segment of the full-length GBV-B clone will be replaced with HCV sequence extending from 5' end of the analogous domain II within the HCV IRES (nt 42) to the initiator codon at the 5' end of the HCV open reading frame (nt 341) to construct the candidate chimera, "GB/C:IRES". The source of HCV cDNA for these studies will be the infectious HCV clone, pCV-H77C, which contains the sequence of the genotype 1a Hutchinson strain virus (Yanagi *et al.*, 1998), whose infectivity in a chimpanzee following intrahepatic inoculation with synthetic RNA transcribed from pCV-H77C has been confirmed.

This GB/C:IRES construct will retain two upstream hairpins within the GBV-B sequence (stem-loops Ia and Ib), and it is thus analogous to the viable "BVDV+HCVdelB2B3H1" chimera of Frolov *et al.* (Frolov *et al.*, 1998). A second chimera can be constructed in which the entire HCV 5' nontranslated RNA will be inserted in lieu of nts 62-446 of the GBV-B genome ("GB/C:5'NTR"). This construct will add to the inserted HCV sequence the most 5' stem-loop

from HCV (stem-loop I). A similar insertion was shown to substantially increase the replication capacity of BVDV+HCVdelB2B3H1 by Frolov *et al.* (Frolov *et al.*, 1998), providing a replication phenotype similar to wild-type BVDV in cell culture.

It is important to point out that there is strong evidence from multiple lines of investigation indicating that it will not be necessary to include coding sequence in these IRES chimeras. This is the case even though Reynolds *et al.* (Reynolds *et al.*, 1995) have argued that the HCV IRES extends past the initiator codon, and into the core-coding region of that virus. Although Lu and Wimmer (Lu and Wimmer, 1996) found it necessary to include HCV core sequence to obtain a viable chimeric poliovirus, the BVDV chimeras reported by Frolov *et al.* (Frolov *et al.*, 1998) did not contain any HCV coding sequence. This discrepancy may be explained by the observation that the only downstream requirement for full activity of both the GBV-B and HCV IRES elements is the presence of an unstructured RNA segment (Honda *et al.*, 1996; Rijnbrand *et al.*, 1999). Presumably, this facilitates interaction of the viral RNA with the 40S ribosome subunit in the early steps of cap-independent translation (Honda *et al.*, 1996). The 5' GBV-B coding sequence fulfills this criterion (Rijnbrand *et al.*, 1999).

EXAMPLE 7

In vitro Characterization of the Translational Activity of IRES Chimeras

The fidelity of the genome-length chimeric constructs will be confirmed by sequencing any DNA segments that have been subjected to PCR during the construction process, as well as confirming sequence at the junction sites. In addition, the translational activity of synthetic RNA derived from these constructs will be assessed and compared to the translational activity of the wild-type GBV-B and HCV RNAs. These studies will be carried out in a cell-free translation assay utilizing rabbit reticulocyte lysates (Rijnbrand *et al.*, 1999). Synthetic RNA will be produced by runoff T7 RNA polymerase transcription using as template ClaI-digested plasmid DNA (BamHI digestion in the case of the genome HCV construct) (T7 Megascript kit, Ambion). ³H-UTP will be added to the reaction mix to allow for quantification of the RNA product. Reticulocyte lysates (Promega) will be programmed for translation by the addition of RNA (at least 50% full-length as determined by agarose gel electrophoresis) at 20, 40 and 80 µg/ml, and translation reactions will be supplemented with microsomal membranes (Promega). ³⁵S-Methionine-labelled translation products will be separated by SDS-PAGE, and the quantity of E1 protein produced from each RNA determined by PhosphorImager analysis (Molecular Dynamics). Comparisons of the activity of the HCV IRES in the background of GBV-B and HCV will take into account differences in the methionine content of the E1 proteins of these

viruses. Based on previous studies of both the GBV-B and HCV IRES elements (Honda *et al.*, 1996; Rijnbrand *et al.*, 1999), it is expected that these studies will confirm that the HCV IRES will retain nearly full activity when placed within the GBV-B background.

EXAMPLE 8

In vivo Characterization of IRES Chimeras

5 Synthetic RNAs produced from each of the two chimeric GBV-B/HCV constructs (GB/C:IRES and GB/C:5'NTR) will be tested for their ability to induce infection and cause liver disease in susceptible tamarins. These studies will be carried out as described in Example 2. GB/C:5'NTR may generate viremia and liver injury more closely resembling that observed with
10 wild-type GBV-B infection (Frolov *et al.*, 1998).

EXAMPLE 9

Chimeric Flaviviruses Containing the HCV NS3 Serine Proteinase/Helicase within the GBV-B Background

Chimeric flaviviruses containing the HCV NS3 serine proteinase/helicase within the
15 GBV-B background are also contemplated within the present invention. The construction of chimeric flaviviruses containing specific heterologous functional polyprotein domains, however, poses a number of special problems. Unlike the situation with the IRES, where the relevant RNA segments appear to have a unique function restricted to cap-independent translation initiation and interact with host cell macromolecules, viral proteins often have multiple functions
20 and may form specific macromolecular complexes with other viral proteins that are essential for virus replication (Lindenbach and Rice, 1999). Furthermore, such chimeric polyproteins must be amenable to efficient processing by the viral proteinases (NS2/NS3 or NS3). This requires knowledge of the proteinase cleavage specificities as well as specific sites of proteolytic cleavage. Although to date there have been no published studies of the processing of the GBV-B
25 polyprotein, the relatively close relationship between GBV-B and HCV, about 30% overall amino acid identity within the polyprotein (Muerhoff *et al.*, 1995), allows good computer predictions of the alignments of these proteins. The crystallographic structures of both the proteinase and helicase domains of the HCV NS3 protein have been solved (Yao *et al.*, 1997). Thus, both linear alignments and models of the 3D structure of the NS3 proteins of these viruses
30 can provide guidelines for designing specific chimeric fusions that are likely to preserve function.

EXAMPLE 10

NS3 Proteinase-Domain Chimeras

In HCV, NS3 contains the major serine proteinase that is responsible for most cleavage events in the processing of the nonstructural proteins, *i.e.*, those that occur at the NS3/4A, 4A/4B, 4B/5A and 5A/5B junctions. The active proteinase domain of HCV is located within the amino terminal third of the NS3 protein (residues 1-181), which shares 31% amino acid identity with the analogous segment of the GBV-B polyprotein (GBV-B vs HCV-BK) (Muerhoff *et al.*, 1995). Importantly, the active site of this proteinase appears to be particularly well conserved in GBV-B. The GBV-B proteinase maintains the residues that are responsible for catalysis and zinc binding in the HCV enzyme (Muerhoff *et al.*, 1995), and unlike the NS3 proteinases of some other flaviviruses preserves the Phe-154 residue that determines in part the S₁ specificity pocket of the enzyme and the preference of the HCV proteinase for substrates with a cysteine residue at the P1 position (Scarselli *et al.*, 1997). Thus, it is not surprising that the relevant proteolytic cleavage sites within the GBV-B polyprotein that are predicted from alignments with the HCV polyprotein all possess a Cys residue at this position. Of greatest significance for the proposed studies, however, is the work of Scarselli *et al.* (Scarselli *et al.*, 1997) who demonstrated that the GBV-B NS3 proteinase is able to effectively process the polyprotein of HCV in studies carried out *in vitro*. Using synthetic peptide substrates, these investigators demonstrated that the enzymatic activities of the GBV-B proteinase (residues 1-181) had kinetic parameters similar to the HCV proteinase on NS4A/4B and NS4B/4A substrates HCV. They did not possess reagents allowing a determination of whether the HCV proteinase is able to cleave a GBV-B substrate, but their results indicate that these viral proteinases share important functional properties. Therefore, these similarities suggest that the HCV proteinase could function in lieu of the GBV-B proteinase if used to replace this segment of an infectious GBV-B clone. In addition, studies with sindbis/HCV chimeras have shown that the HCV proteinase can cleave within the framework of a sindbis polyprotein (Filocamo *et al.*, 1997).

In considering the design of these NS3 proteinase chimeras, there are two additional important considerations. First, in HCV, the cleavage between NS2 and NS3 occurs *in cis*, as the result of a zinc-dependent metalloproteinase that spans the NS2/NS3 junction (Hijikata *et al.*, 1993). As only the NS3 sequences will initially be exchanged, the viability of the resulting chimeras will be dependent upon preservation of the *cis*-active cleavage across a chimeric NS2/NS3 proteinase domain. The alignment of GBV-B and HCV sequences shows that residues in HCV that have been shown by Grakoui *et al.*, 1993, to be essential for the NS2/NS3 cleavage

are conserved in GBV-B (Muerhoff *et al.*, 1995). Additional chimeras that will include the relevant carboxyl-terminal portion of NS2 can also be created.

A second important consideration is that the mature HCV NS3 proteinase functions as a noncovalent assembly of the NS3 proteinase domain and the amino terminal portion of NS4A, a proteinase accessory factor. The details of this association are well known, and have been studied at the crystallographic level (Kim *et al.*, 1996). The N-terminal domain of the folded proteinase contains eight β strands, including one contributed directly by the NS4A peptide backbone. X-ray studies have shown that this array of β strands gives rise to a much more ordered N-terminus. Thus, the presence of the NS4A strand seems likely to contribute to the structure of the substrate-binding pocket. It is not known whether the NS3 proteinase of GBV-B also requires a similar interaction with NS4A of that virus for complete activity, or, if so, whether the NS4A of GBV-B could substitute for NS4A of HCV in forming the fully active NS3 proteinase of HCV. The predicted GBV-B NS4A molecule is 54 amino acid residues in length (Simons, *et al.*, 1995; Muerhoff *et al.*, 1995), just as in HCV. However, the level of amino acid homology between the NS4A molecules is not especially high, and the potential interaction with either NS3 molecule cannot be predicted from this sequence on the basis of available knowledge. To overcome this potential problem, chimeras will be created in which not only the NS3 proteinase domain of GBV-B is replaced, but also the relevant NS4A segment as well, with homologous segments of the HCV polyprotein. The interaction of the HCV NS3 and NS4A domains represents a unique target for antiviral drug design, and it would be beneficial to have this specific interaction present in any virus to be used as a surrogate for HCV in the evaluation of candidate antiviral inhibitors of HCV proteinase *in vivo*.

The NS3 proteinase chimeras that can be made include "GB/C:NS3^P", which will contain the sequence encoding the first 181 amino acid residues of the HCV NS3 molecule in lieu of that encoding the first 181 residues of GBV-B NS3, and "GB/C:NS3^P4A", which will include the same NS3 substitution as well as the HCV sequence encoding the amino-terminal segment of NS4A that forms the interaction with NS3. The precise NS4A sequence to be included in the latter chimera will be based on the modeling studies, which may also suggest more effective fusions of the NS3 proteinase domain of HCV with the downstream NS3 helicase domain of GBV-B. The source of HCV cDNA for these studies will be the infectious HCV clone, pCV-H77C, which contains the sequence of the genotype 1a Hutchinson strain virus (Yanagi *et al.*, 1998).

EXAMPLE 11**NS3 Helicase Domain Chimeras**

In addition to serine proteinase activity located within the amino-third of NS3, the downstream carboxy-terminal two-thirds of the molecule contains an RNA helicase activity.

5 These two functional domains appear to be separated by a flexible spacer, within which the fusion of HCV proteinase or helicase domain sequences with GBV-B sequence will be placed. The exact role of the helicase in the HCV life-cycle is not known, but it is almost certainly required for dsRNA strand-separation during some phase of viral RNA synthesis. The helicase domains of GBV-B and HCV are remarkably well conserved, with some regions within the
10 helicase showing as much as 55% amino acid identity (Muerhoff *et al.*, 1995). The GBV-B helicase is more closely related to the HCV helicase than all other flaviviral NS3 helicases, and it preserves many residues found within the conserved helicase motifs of HCV. Thus the HCV NS3 helicase may be capable of functioning when placed within the polyprotein of GBV-B, and such a chimeric virus may be capable of replication. Residues 182-620 of the GBV-B NS3
15 molecule will be substituted with the analogous segment of HCV (FIG. 5, "GB/C:NS3^h"). A chimera will also be made in which the entire NS3 and amino terminal NS4 protein sequences of GBV-B is replaced with the homologous HCV sequences ("GB/C:NS3-4A"). The latter construct will thus represent a dual proteinase-helicase chimera. As with the proteinase chimeras, the HCV cDNA will be derived from pCV-H77C (Yanagi *et al.*, 1998).

EXAMPLE 12***In vitro* Characterization of NS3 and NS3-NS4A Chimera**

Prior to being evaluated for infectivity in susceptible tamarins, RNAs produced *in vitro* from these clones will be characterized *in vitro*. This evaluation will be restricted to a documentation of the proper processing of the expressed polyprotein (*i.e.*, NS2/NS3 and NS3
25 proteinase functions), since there are no relevant assays that can determine whether the helicase or RNA-dependent RNA polymerase activities in these polyproteins are sufficient for virus replication. The proteolytic processing of the polyprotein is important, however, as it may be altered either by inclusion of the heterologous HCV NS3 proteinase in lieu of the natural GBV-B protease, or by a change in the folding of the polyprotein induced by inclusion of HCV sequence
30 anywhere within the polyprotein. These studies will be carried out in cell-free coupled transcription/translation assays ("TnT" system, Promega) supplemented with microsomal membranes. Template DNAs will be digested with Sal I, which restricts the cDNA within the NS5B coding region. ³⁵S-methionine-labelled translation products will be separated by SDS-

PAGE, and the mature NS3 protein identified by its apparent molecular mass. The NS3 and NS5B proteins will be identified by immunoblot analysis using rabbit antisera to the GBV-B NS3 and NS5B proteins. Generation of a mature ~68 kDa NS3 protein will provide proof of both the *cis*-active NS2/NS3 cleavage and the NS3-mediated cleavage of NS3/NS4A. Similarly, identification of a mature, processed NS5B molecule will provide further support for the activity of the NS3 proteinase. Controls for these studies will be the wild-type GBV-B polyprotein expressed in similar fashion from the full-length GBV-B clone. If necessary to more clearly demonstrate the processing of the nonstructural proteins in these constructs, subclones representing the nonstructural region of the chimeric sequences could be produced.

EXAMPLE 13

In vivo Characterization of NS3 and NS3-4A Chimeras

Synthetic RNAs produced from each of the chimeric GBV-B/HCV constructs described in the preceding section will be tested for their ability to induce infection and cause liver disease in susceptible tamarins. These studies will be carried out using the approach described above.

EXAMPLE 14

Chimeric Flaviviruses Containing the HCV NS5B RNA Dependent RNA Replicase within the GBV-B Background

The HCV NS5B molecule contains an RNA-dependent RNA polymerase that plays a central role in replication of the virus. Although this molecule represents a prime target for drug discovery efforts, it has proven difficult to express NS5B in a form that retains enzymatic activity specific for HCV RNA as a substrate. Thus, relatively little is known of the functional activity of the HCV replicase, including structure-function relationships of NS5B. Despite this, the NS5B proteins of GBV-B and HCV appear to be functionally closely related, as they share as much as 43% amino acid identity (Muerhoff *et al.*, 1995). A more important question may be whether an RNA dependent RNA polymerase can act on foreign substrates. However, published work has shown that *in vitro* purified HCV polymerase has very little specificity for its template, using hepatitis C or globin message with equal fidelity (Behrens *et al.*, 1996; Al *et al.*, 1998). This finding is very similar to that obtained with picornaviral polymerases, where it has been known for many years that *in vitro* the enzyme exhibits very little specificity. It has always been considered highly likely that this situation would not pertain *in vivo* where it was thought that the interaction of viral or cellular factors with the 3' end of the genome would generate template specificity. However, recent reports have shown that the removal of the entire 3' untranslated

sequence (leaving, however, the poly(A) region present) from both the poliovirus and rhinovirus genome does not completely abrogate the infectivity of the virus (Todd *et al.*, 1997). Furthermore, virus, which was recovered after the initial transfections, was shown to have recovered much of the infectivity of the original virus (Todd *et al.*, 1997). The mechanism for this recovery of infectivity is at present unknown, but these results suggest that the HCV polymerase may be able to function to replicate infectious GBV-B/HCV NS5B chimeras.

Thus a chimeric genome-length virus can be created in which the NS5B coding sequence of HCV (amino acids 2422-3014, 593 residues) is inserted within the background of GBV-B in lieu of its native RNA-dependent RNA polymerase (amino acids 2274-2864, 591 residues). This chimeric virus would be valuable for animal studies of candidate antiviral inhibitors of HCV RNA synthesis.

This NS5B chimera would be evaluated to determine that there was proper proteolytic processing of the polyprotein. This would be accomplished by expression of the chimeric polyprotein in a coupled translation-transcription reaction, followed by immunoblot analysis for the mature NS5B protein, as described for the NS3 and NS3-4A chimeras in the preceding section. If these results confirm that the GB/C:5B chimeric polyprotein is processed with release of NS5B, studies in tamarins would progress to determine whether synthetic RNA transcribed from the clone is infectious and capable of causing liver disease in intrahepatically inoculated animals. These studies would be carried out as described above.

A chimeric molecule can be constructed from an infectious GBV-B clone in which the HCV NS3 proteinase or proteinase/helicase sequence would be placed in frame in lieu of the homologous GBV-B sequence, and this chimeric cDNA would be used to generate infectious GBV-B/HCV chimeric viruses by intrahepatic inoculation of synthetic RNA in tamarins. Published studies indicate that the GBV-B and HCV proteinases have closely related substrate recognition and cleavage properties, likely making such chimeras viable and capable of initiating viral replication in appropriate cell types.

EXAMPLE 15

Chimeric Viruses Containing HCV Structural Proteins within a GBV-B Genetic Background, and GBV-B Structural Proteins within an HCV Background

It is well documented that the structural proteins of one flavivirus may in some cases be substituted for those from another member of the family. Such chimeric viruses have been recovered from viruses as distantly related to each other as dengue virus and tick-borne encephalitis virus (Pletnev *et al.*, 1992). More recently, the prM and E proteins of Japanese

encephalitis virus have been used to replace the equivalent proteins in a vaccine strain of yellow fever virus to produce a JE/YF chimera (Chambers *et al.*, 1999). These observations suggest that chimeras in which the structural proteins of HCV have been used to replace the homologous proteins of GBV-B may well be viable and capable of replication. The isolation of a chimeric virus containing HCV structural proteins, but having the growth characteristics of GBV-B virus, could answer many fundamental questions concerning the structure and interaction of these proteins in HCV. They would also be useful in addressing the nature of the immune response to HCV structural proteins in infected primates (Farci *et al.*, 1992). More to the point of this application, the availability of such chimeric viruses would allow studies of candidate HCV vaccines to be carried out in the tamarin model. This would be a major advance, because at present such studies are limited to chimpanzees (Choo *et al.*, 1994).

The basis for the difference in the host ranges of HCV and GBV-B is completely unknown. Among many other possibilities, it is conceivable that the host range is dependent upon the availability of a specific receptor(s). If this were the case, host range might be dependent upon the envelope proteins that must interact with the putative cellular receptor. Thus, a chimeric virus containing the envelope proteins of HCV within the genetic background of GBV-B might be noninfectious in tamarins (but potentially infectious in chimpanzees). Thus, a finding that both structural protein chimeras are noninfectious in the tamarin, may require the construction of complementary chimeras in which the relevant GBV-B structural proteins will be inserted into the background of an infectious HCV clone. If inclusion of the GBV-B envelope proteins within the backbone of HCV confers on the resulting chimera the ability to replicate in tamarins, it will confirm an important role for the structural proteins in defining the different host ranges of these viruses. More importantly, the resulting virus would be an exceptionally valuable resource for future studies as it would contain all of the nonstructural replication elements, as well as the 5' and 3' nontranslated regions, of HCV. Such a virus would allow the tamarin model to be used to address many unresolved issues in HCV biology and pathogenesis.

EXAMPLE 16

Construction and Evaluation of Structural Protein Chimeras

In designing structural protein chimeras, it is important to note that the two envelope proteins of HCV, E1 and E2, form noncovalent heterodimeric complexes that are likely to be important in the assembly of infectious virus particles. This is not known to be the case with the envelope proteins of GBV-B, but it is likely given similarities in the sizes and hydropathy profiles of these proteins (Simons *et al.*, 1995; Muerhoff *et al.*, 1995). Accordingly, the E1 and

E2 proteins will be replaced as a unit, and chimeras containing only one of these proteins from the heterologous virus will generally not be produced. First, a chimera will be created where the E1 and E2 regions of GBV-B virus are replaced with those of HCV, "GB/C:E1-2". The source of HCV cDNA for these constructions will be pCV-H77C (Yanagi *et al.*, 1998). A chimera will also be made in which the core protein, in addition to the envelope proteins, is replaced with the homologous proteins of HCV ("GB/C:Co-E2"). Additional chimeras will be made to determine whether tamarins can be infected with chimeras containing the GBV-B structural proteins within the genetic background of HCV. These will include "C/GB:E1-2" and "C/GB:Co-E2". The backbone for these chimeras will be pCV-H77C, the infectious genotype 1a cDNA clone developed in the Purcell laboratory at NIAID (Yanagi *et al.*, 1998).

The specific amino acid sequences of GBV-B to be replaced with the homologous segments of HCV have been determined by alignments of the GBV-B and HCV sequences, coupled with the location of signalase cleavage sites predicted to be present within the amino terminal third of the GBV-B polyprotein using the computer algorithm of Von Heijne. These predicted signalase cleavages lie between residues 156/157 (core/E1), aa 348-349 (E1/E2) and 732/733 (E2/NS2) in the GBV-B sequence. Thus, the chimera GB/C:E1-2 will contain sequence encoding HCV aa 192-809 in lieu of that encoding aa 157-732 in GBV-B, while the insertion in the GB/C:Co-E2 chimera will extend from the initiator AUG codon (aa 1) to residue 809 in HCV, and will be spliced into GBV-B in lieu of the segment encoding aa 1-732 in the GBV-B clone. The complementary chimeras to be constructed within the background of HCV will involve exchanges of the same segments of the genomes.

EXAMPLE 17

Characterization of Structural Protein Chimeras

Prior to being evaluated for infectivity in tamarins, the processing of these chimeric polyproteins will be examined in coupled transcription/translation reactions supplemented with microsomal membranes, as described in the preceding sections for the proteinase and other proposed chimeras. If these results confirm that the polyprotein is processed as expected, with production of glycosylated E1 and E2 proteins from each of the chimeras (and seen in similar studies with HCV proteins), studies would proceed in tamarins as previously described. The results of these studies may provide novel information on the basis of the host range differences that exist between HCV and GBV-B. If these results suggest that the envelope proteins play a critical role in determining host range, additional studies could be carried out with these chimeras in chimpanzees (which are permissive for HCV but apparently not for GBV-B).

EXAMPLE 18**Further Characterization of Rescued Chimeric Viruses**

Where infection with chimeric viruses is induced in animals that are injected within the liver with synthetic RNA, this virus will be passaged in GBV-B naïve tamarins to further
5 characterize the nature of the infection induced by the chimera. This will be accomplished by taking a pool of the 3 highest titer GBV-B RNA-containing serum specimens from the animal that was successfully transfected with RNA, and inoculating 1 mL of a 1:100 dilution of this pool intravenously into two susceptible animals. These animals will be monitored for infection and
10 liver disease. These animals will be followed until resolution of the viremia and appearance of antibodies detectable in immunoblots with GST-NS3 protein expressed in *E. coli*, or for at least 6 months should an animal sustain a chronic infection. RT-PCR amplification of chimeric segments of the genome may be employed to determine whether the altered phenotype results from mutations within the heterologous portion of the genome.

EXAMPLE 19**Use of GBV-B as Model for HCV**

GBV-B and/or GBV-B/HCV chimeras can be used as a model for HCV. Such studies will allow one to investigate the mechanisms for the different biological properties of these viruses and to discover and investigate potential inhibitors of specific HCV activities (*e.g.*,
20 proteinase) required for HCV replication.

EXAMPLE 20**Use of GBV-B/HCV Chimeras to Test Candidate HCV NS3
Proteinase Inhibitors or Other Inhibitors of HCV**

GBV-B/HCV viruses can be used in preclinical testing of candidate HCV NS3 proteinase
25 inhibitors or other inhibitors of HCV.

1. Candidate Substances

As used herein the term "candidate substance" refers to any molecule that is capable of
30 modulating HCV NS3 proteinase activity or any other activity related to HCV infection. The candidate substance may be a protein or fragment thereof, a small molecule inhibitor, or even a nucleic acid molecule. It may prove to be the case that the most useful pharmacological compounds for identification through application of the screening assay will be compounds that are structurally related to other known modulators of HCV NS3 proteinase activity. The active

compounds may include fragments or parts of naturally-occurring compounds or may be only found as active combinations of known compounds that are otherwise inactive. However, prior to testing of such compounds in humans or animal models, it will be necessary to test a variety of candidates to determine which ones have potential.

5 Accordingly, the active compounds may include fragments or parts of naturally-occurring compounds or may be found as active combinations of known compounds that are otherwise inactive. As such, the present invention provides screening assays to identify agents that are capable of inhibiting proteinase activity in a cell infected with chimeric GBV-B/HCV viruses containing the HCV proteinase. It is proposed that compounds isolated from natural sources,
10 such as animals, bacteria, fungi, plant sources, including leaves and bark, and marine samples may be assayed as candidates for the presence of potentially useful pharmaceutical agents. It will be understood that the pharmaceutical agents to be screened could also be derived or synthesized from chemical compositions or man-made compounds. Thus, it is understood that the candidate substance identified by the present invention may be polypeptide, polynucleotide, small molecule
15 inhibitors or any other compounds that may be designed through rational drug design starting from known inhibitors of proteinases or from structural studies of the HCV proteinase.

 The candidate screening assays are simple to set up and perform. Thus, in assaying for a candidate substance, after obtaining a chimeric GBV-B/HCV virus with infectious properties, a candidate substance can be incubated with cells infected with the virus, under conditions that
20 would allow measurable changes in infection by the virus to occur. In this fashion, one can measure the ability of the candidate substance to prevent or inhibit viral replication, in relationship to the replication ability of the virus in the absence of the candidate substance. In this fashion, the ability of the candidate inhibitory substance to reduce, abolish, or otherwise diminish viral infection may be determined.

25 “Effective amounts” in certain circumstances are those amounts effective to reproducibly reduce infection by the virus in comparison to the normal infection level. Compounds that achieve significant appropriate changes in activity will be used. Candidate compounds can be administered by any of a wide variety of routes, such as intravenously, intraperitoneally, intramuscularly, orally, or any other route typically employed.

30 It will, of course, be understood that all the screening methods of the present invention are useful in themselves notwithstanding the fact that effective candidates may not be found. The invention provides methods of screening for such candidates, not solely methods of finding them.

2. *In vitro* Assays

In one particular embodiment, the invention encompasses *in vitro* screening of candidate substances. Using a cell line that can propagate GBV-B in culture, *in vitro* screening can be used such that GBV-B or HCV virus production or some indicator of viremia is monitored in the presence of candidate compounds. A comparison between the absence and presence of the candidate can identify compounds with possible preventative and therapeutic value.

3. *In vivo* Assays

The present invention also encompasses the use of various animal models to test for the ability of candidate substances to inhibit infection by HCV. This form of testing may be done in tamarins.

The assays previously described could be extended to whole animal studies in which the chimeric virus could be used to infect a GBV-B permissive primate, such as a tamarin. One would then look for suppression of viral replication in the animal, and a possible impact on liver disease related to replication of the infectious chimeric virus. The advantage of this *in vivo* assay over present available assays utilizing HCV infection in chimpanzees is the reduced cost and greater availability of GBV-B permissive nonhuman primate species.

Treatment of animals with test compounds will involve the administration of the compound, in an appropriate form, to the animal. Administration will be by any route that could be utilized for clinical or non-clinical purposes, including but not limited to oral, nasal, buccal, rectal, vaginal or topical. Alternatively, administration may be by intratracheal instillation, bronchial instillation, intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection. Specifically contemplated are systemic intravenous injection, intraperitoneal injection, and oral administration.

Determining the effectiveness of a compound *in vivo* may involve a variety of different criteria. Such criteria include, but are not limited to, survival, reduction of rate of infection, arrest or slowing of infection, elimination of infection, increased activity level, improvement in liver function, and improved food intake.

EXAMPLE 21

Use of Infectious GBV-B/HCV Chimeras as Vaccines

Infectious GBV-B/HCV chimeras expressing HCV envelope proteins will have utility as a vaccine immunogen for hepatitis C. Such clones clearly have the potential to be constructed as

chimeras including relevant hepatitis C virus sequences in lieu of the homologous GBV-B sequence, providing unique tools for drug discovery efforts.

Chimeric viruses containing the envelope proteins of hepatitis C virus would confer the antigenic characteristics of hepatitis C virus on the chimera. These chimeras may have the ability to replicate in chimpanzees (and thus humans) by virtue of the fact that the chimeric envelope is now able to interact with the human hepatocyte cell surface, a necessary first step in virus replication. Therefore, the chimeric virus, while able to infect and replicate in humans, may not cause much or any disease-the reasoning here is that the genetic backbone of the chimera that encodes the nonstructural proteins of GBV-B has not evolved for replication in human cells and thus may not replicate well. Thus, the chimera may have limited replication ability, cause no disease, but still elicit immunity to the surface envelope proteins of HCV and thus have potential as a hepatitis C vaccine. These chimeras can be tested for their ability to promote immunity to HCV through an immune response.

EXAMPLE 22

Construction of a GBV-B Infectious Clone

A genome-length cDNA copy of the complete GB virus B (GBV-B) genome sequence, including the novel 3' terminal sequence, was assembled from fragments amplified by reverse transcription-polymerase chain reaction (RT-PCR) from viral RNA present in a 0.2 µl aliquot of infectious serum (GB agent pool mystrax 666, 8/93) supplied by Dr. Jens Bukh of the National Institutes of Health. The serum sample was diluted with 100 µl of fetal calf serum and extracted using the Trizol system (GIBCO/BRL). The pellet was dissolved in 10 mM dithiothreitol containing 20 u/mL RNasin (Promega). The selection of primers for cDNA synthesis and PCR amplification was based on the published sequence of GBV-B (Simons *et al.*, 1995). RT/PCR was carried out using Superscript Reverse Transcriptase (GIBCO/BRL) and the Advantage cDNA PCR Kit (Clontech). Four subgenomic regions were amplified covering the entire published GBV-B genome sequence. A fifth subgenomic region included the novel 3' terminal sequence that was identified in our laboratory. The oligonucleotide primer sets (listed 5'→3') used for amplification of the individual regions included:

Primer 1: 5'CGGGATCCCGTAATACGACTCACTATAGACCACAAACACTCCAGT TTG –
3' (SEQ ID NO:3)

Primer 2: 5'-GTGGAATTCACAGCGTCATA-3' (SEQ ID NO:4) (Places a T7 RNA polymerase promoter sequence immediately upstream of the GBV-B sequence; the amplified

segment extends from the 5' terminus of the viral genome to the unique *EcoRI* site at position 1978.)

Primer 3: 5'-TGTGAATTCCACTCTCCTACC-3' (SEQ ID NO:5)

Primer 4: 5'-TTATCGATTGCAGCAACCATG-3' (SEQ ID NO:6) (Overlapping *EcoRI* site at position 1978 and the unique *ClaI* site at 5327.)

Primer 5: 5'-CATGGTTGCTGCAATCGATAAGCTGAAGAGTACAATAAC-3' (SEQ ID NO:7)

Primer 6: 5'-GACAACAGACGCTTGACACG-3' (SEQ ID NO:8) (Overlapping the *Cla I* site at 5327 and a unique *Sal I* site at 7847 in the published sequence: however, the *Sal I* site was not present in the amplified GBV-B sequence, and was thus introduced into the cDNA using the QuikChange Site-directed Mutagenesis Kit (Stratagene).)

Primer 7: 5'-CTGTCATGGGAGATGCGTAC-3' (SEQ ID NO:9)

Primer 8: 5'-CGAGCTCGAGCACATCGCGGGGTCGTTAAGCCCCGGGGTCTCC-3' (SEQ ID NO:10) (Overlapping the *Sal I* site at 7847 and the published 3' end of the genome.)

Primer 9: 5'-GACAACAGACGCTTGACACG-3' (SEQ ID NO:11)

Primer 10: 5'-CCGACTCGAGAATTCGGCCCTGCAGGCCACAACAGTCTCGCGAG TTTTAAATTCCAAGCGGGGGTTGCCCTCCGCTTGGAACAAAAACCGGGGTGCAGCCC TTGGTAC-3' (SEQ ID NO:12) (Overlaps product of primers 7 and 8 and extends to the novel 3' terminal sequence; includes an *Xho I* site at the 3' terminus of the GBV-B sequence to aid subsequent manipulations.)

RT-PCR products were gel-purified using the QIAquick Gel Extraction Kit (Qiagen) and ligated into a plasmid vector with the PstBlue-1 Perfectly Blunt Cloning Kit (Novagene). Bacterial colonies were screened for plasmid DNAs with the correct insert size. Those harboring an insert of the approximate correct size were subjected to diagnostic restriction analysis. cDNA inserts generating the appropriate restriction patterns were sequenced in both directions using an ABI DNA sequencer, and those with sequences closest to the published GBV-B sequence (Simons *et al.*, 1995) were selected for subsequent assembly into the full-length clone.

The cloned overlapping cDNA fragments were assembled into the pACNR1180 plasmid using the unique restriction sites noted in the primer descriptions. The final, fully assembled plasmid bearing a 9.4 kb cDNA GBV-B insertion was subjected to DNA sequencing to confirm the validity of the construction and the absence of any mutations or errors in the sequence that may have been introduced during the assembly process. The sequence of the GBV-B infectious cDNA is shown in SEQ ID NO:2.

Plasmid DNA containing the GBV-B cDNA was purified by isopycnic centrifugation on a CsCl gradient and isolated by ethanol precipitation. The plasmid DNA was linearized at the 3' terminus of the cDNA insert by digestion with *Xho* I. Two µg of the linearized DNA was used as template in an *in vitro* transcription reaction using the Ambion T7 MEGAscript Kit according to the manufacturer's instructions. The integrity of the resultant synthetic RNA product was confirmed by nondenaturing agarose gel electrophoresis.

Approximately 60 µg of RNA transcribed from the GBV-B cDNA clone was diluted in phosphate buffered saline and injected under direct visualization at laparotomy into the liver of a healthy tamarin without a prior history of GBV-B infection (*Saguinus oedipus*). Serum samples were collected from the animal weekly beginning at 2 weeks post-inoculation and tested for the presence of GBV-B RNA by real-time quantitative RT-PCR (TaqMan® assay). Results are shown in Table 2.

Table 2. Serum samples collected from the animal weekly beginning at 2 weeks post-inoculation and tested for the presence of GBV-B RNA by real-time quantitative RT-PCR.

Week Postinoculation	Genome equivalents/ml
0	0
2	8.3×10^3
3	6.6×10^5
4	4.6×10^8

The presence of high and increasing titers of viral RNA in the serum of this animal between 2-4 weeks postinoculation confirms the infectivity of the full-length cDNA clone containing the novel 3' terminal sequence. The presence of viremia demonstrated that viral replication had ensued following the delivery of the synthetic viral RNA to hepatocytes.

EXAMPLE 23

Construction of a Chimeric Virus Containing the Domain III of the 5' NTR of HCV (IRES) Within the Genetic Background of GBV-B.

1. Construction of plasmids containing chimeric cDNAs.

As described herein, a genome-length molecular clone of GBV-B was assembled from a series of overlapping cDNA fragments produced by RT-PCR from viral RNA isolated in a GBV-B-infected tamarin serum. This clone incorporates the novel GBV-B 3'NTR sequence, as described herein. The cDNA has been placed downstream of a T7 RNA polymerase promoter in the context of the bacterial vector pACNR1180. To facilitate further cloning steps, a C → T substitution was introduced at nucleotide (nt) position 496 of the GBV-B cDNA, so that a unique

MluI restriction site was generated in the core-coding sequence without modifying the amino acid sequence. The resulting plasmid containing the full-length GBV-B cDNA with the engineered MluI site at nt position 491 will be referred to as pGBV-B/Mlu. Chimeric GBV-B / HCV 5'NTRs were generated by substituting domains of the GBV-B 5'NTR by HCV counterparts using a gene fusion PCR strategy (Landt *et al.*, 1990) and HCV sequences derived from Hutchinson strain of genotype 1a, Genbank accession numbers AF011751 (SEQ ID NO:13), AF011752, and AF011753, each of which is incorporated herein by reference (Inchauspe *et al.*, 1991). The invention is not limited to a particular strain of HCV and sequences found in other HCV sequences may also be used, such as Genbank accession numbers AF009606, M67463, AF290978, AF387806, AF271632, M62321, AF387807, AF387805, AF387808, D10749, AF511948, AF177040, AF177038, AF177039, AF177037, AF511949, or M32084, each of which is incorporated herein by reference. Chimeric 5'NTRs were amplified using overlapping PCR fragments spanning the desired HCV region framed by appropriate GBV-B regions, so that the BamHI restriction site located upstream of the T7 promoter sequence and the MluI restriction site (nt 491) were used to substitute the parent GBV-B fragment within the molecular infectious clone. The sequence of all PCR-amplified fragments within the resulting, chimeric pGB/I-II-III^{HC}, pGB/ Δ Ib/II-III^{HC}, pGB/II-III^{HC}, and pGB/III^{HC} plasmids (FIG. 2) were confirmed.

2. *In vitro* and *ex vivo* analysis of translation efficiency of chimeric IRES elements.

To assess the translational efficiencies of RNAs transcribed from the plasmids shown in FIG. 2, some of which contain chimeric IRES viral sequences, translation initiation activities were determined *in vitro* in rabbit reticulocyte lysates and *ex vivo* in primary tamarin hepatocytes. To facilitate quantitation *ex vivo*, the MluI-XhoI fragment (nts 491-9398), spanning almost the entire GBV-B coding sequence and the 3' NTR of the cDNA in pGBV-B/Mlu, was replaced by a fragment encoding Renilla luciferase (RLuc) followed by the GBV-B 3' NTR. This substitution was carried out downstream of the chimeric sequences, generating plasmids pI-II-III^{HC}-Luc, p Δ Ib/II-III^{HC}-Luc, pII-III^{HC}-Luc, and pIII^{HC}-Luc that were used in translation studies.

These plasmids were linearized at the Xho I restriction site and used as templates for *in vitro* transcription using T7 RNA polymerase (Megascript kit, Ambion). *In vitro* translations were performed in Flexi Rabbit Reticulocyte lysates as described by the supplier (Promega) with 125mM KCl in the presence of ³⁵S-Methionine. The RNA transcript concentration was 12.5ng/ μ l, well below the saturation point. Translation products were analyzed by 8% SDS-

PAGE and quantitated by PhosphorImager analysis (Molecular Dynamics). Exemplary results are shown in FIG. 3.

Reporter RNAs representing I-II-III^{HC}, Δ Ib/II-III^{HC}, and II-III^{HC}, all of which contain the complete HCV IRES element, translated with about 2.5-fold higher efficiency as compared to GBV-B/Mlu RNA which contains the intact GBV-B IRES (FIG. 2). The translational activity observed for III^{HC} was similar to that observed for the GBV-B/Mlu RNA. This is remarkable as this RNA contains a chimeric IRES in which upstream and downstream sequences flanking domain III of the HCV IRES are derived from the GBV-B IRES. These data suggest that the fusion of HCV and GBV-B 5'NTR sequences did not adversely affect translational activity of the internal ribosome entry site (IRES) when assayed in the cell-free translation system.

To assess the translational activity of these chimeric reporter RNA transcripts in living tamarin cells, *in vitro*-transcribed RNA was mixed with the lipid-based compound DMRIE (Life Technologies) and transfected into primary tamarin hepatocytes (*S. oedipus*) that were prepared as described previously (Beames *et al.*, 2000, incorporated herein by reference). As an internal control, an RNA transcript encoding firefly luciferase (FLuc) under translational control of the EMCV IRES was cotransfected at 1:19th the abundance of the RLuc reporter constructs.

Translational activity was quantitated by determining the levels of luciferase activity 24 hours following transfection (FIG. 4). In this system, which may represent more closely the conditions present during GBV-B infection of the tamarin, the translational activity of each of the RNA transcripts containing the complete HCV IRES was lower than that observed with the GBV-B/Mlu control. Thus, the GBV-B IRES may be more active in tamarin cells (the normal host species for GBV-B) than the HCV IRES. However, the chimeric transcript III^{HC} which contains domain III of the HCV IRES in lieu of its GBV-B counterpart had translational activity comparable to that of GBV-B/Mlu.

Although the data shown in FIG. 3 and FIG. 4 indicate that there may be variation in the relative activities of these chimeric RNAs in different translation systems (in tamarin hepatocyte cultures and in a cell free translation system prepared from rabbit reticulocytes), these results may indicate that each of the chimeric IRES elements in these RNAs retains substantial translation-initiation activity.

EXAMPLE 24**Infectivity of Chimeric Viral cDNAs Containing HCV 5'NTR Sequences within the Genetic Background of GBV-B.****1. Inoculation of synthetic RNA into GBV-B susceptible tamarins.**

5 After linearization of the plasmid DNAs shown in FIG. 2 by digestion with XhoI, genome-length RNA transcripts containing the chimeric 5'NTR sequences were transcribed from the plasmids using the T7 MegaScript kit (Ambion). An aliquot of the reaction products was tested by agarose gel electrophoresis to ensure RNA integrity and to approximate the quantity of RNA. The remaining RNA was frozen at -80°C until inoculation, without further purification,
10 into the liver of susceptible tamarins. Individual animals (*S. mystax*) received a dose of approximately 100-200 µg of RNA transcribed from only one of the plasmids shown in FIG. 2. The RNAs were inoculated into the liver under direct visualization following exposure by laparotomy. Inoculated animals were followed as described in the preceding sections of this document, with periodic testing of serum ALT, antibody to the NS3 protein of GBV-B, GBV-B
15 genome, and with liver biopsy to assess pathologic changes in the liver. All tamarins were housed at the Southwest Regional Primate Research Center at the Southwest Foundation for Biomedical Research. Animals were cared for by members of the Department of Laboratory Animal Medicine at the Southwest Foundation for Biomedical Research in accordance with the Guide for the Care and Use of laboratory Animals. All protocols were approved by the
20 Institutional Animal Care and Use Committee.

The primary indicator of successful infection was detection of sustained viremia in samples of serum collected over a period of weeks. Viremia was monitored by detection of GBV-B RNA using a sensitive and specific real-time RT-PCR assay. RNA was isolated from virus present in tamarin serum using the QiaAmp viral RNA extraction kit (Qiagen) and
25 amplified in a one-step RT-PCR reaction utilizing TaqMan® EZ core reagents (Perkin Elmer). First-strand cDNA synthesis was carried out using the primer NS5ARPp (5'-GAAGGAGGGAGGTTTGAAGGA-3', position 6949-6969)(SEQ ID NO:14). The cDNA product was quantified in a 5' exonuclease PCR (TaqMan®) assay using a primer-probe combination that recognized sequences in the GBV-B NS5A gene. The primers NS5ARPp and
30 NS5AFPp (5'-CCAGTTCCGGGCAAGAACT-3', position 6844-6862)(SEQ ID NO:15) and probe (nts 6913-6938) were obtained from Fisher and Perkin Elmer, respectively. A synthetic RNA derived from the infectious clone was used as the standard for quantitation of the RNA.

There was no evidence for replication of the I-II-III^{HC}, II-III^{HC}, and Δ Ib/II-III^{HC} chimeras following their inoculation into the liver of individual GBV-B naïve tamarins during the 22-week follow up period. In contrast, RT-PCR suggested the possible presence of a low level of viral RNA in the serum of the animal injected with III^{HC} chimera 1-2 weeks postinoculation, followed by disappearance of any sign of viremia. However, 12 weeks after inoculation with the III^{HC} RNA, viremia reappeared in tamarin T16444, rapidly reaching a titer in excess of 10^7 genome copies per ml, and subsequently persisting for 20 weeks (terminating 32 weeks following inoculation of the RNA) (FIG. 5). The course of the infection in this animal from week 12 on was similar to what is seen in infection of naïve animals with the wild-type infectious clone, with the exception that there was no detectable evidence of antibodies to the NS3 protein. The long delay to the appearance of the secondary viremia in this animal suggests a requirement for the accumulation of adaptive mutations that are compensatory to changes in the replication capacity of the RNA related to the presence of the chimeric 5'NTR sequence.

The results shown in FIG. 5 indicate the replication competence of the III^{HC} chimeric RNA (see FIG. 2) following intrahepatic inoculation of synthetic RNA in a GBV-B naïve tamarin (*S. mystax*). The presence of the chimeric 5'NTR sequence was confirmed in the RNA replicating in this animal by RT-PCR amplification of RNA extracted from the serum, followed by direct DNA sequencing of the amplified cDNA.

To prove that inoculation of T16444 with synthetic III^{HC} chimeric RNA had led to the rescue and replication of virus containing the chimeric 5'NTR sequence, serum collected 14 weeks after the inoculation of this animal with RNA was used to inoculate a second GBV-B naïve animal (T16451) by intravenous injection. This is a stringent test for the presence of virus, since nonpackaged viral RNA that might be released from the liver into the serum would be expected to be highly degraded and no longer infectious. Only viral RNA packaged into viral particles would be expected to transmit the infection. This second animal rapidly developed evidence of GBV-B infection, with viremia detected as early as 1 week after infection, and with a peak viremia of greater than 10^8 genome copies/ml by 4 weeks postinfection. The rapid appearance of the viremia and early robust replication of the virus in this second animal suggested that the chimeric virus had indeed undergone adaptation and that it no longer required a lengthy incubation period to induce viremia.

These results confirm that the synthetic chimeric III^{HC} RNA replicated and produced infectious virus particles, capable of transmitting the infection to a second animal. Liver biopsy in this animal confirmed the presence of hepatic inflammation, while serial determinations of

serum ALT activities demonstrated the typical profile of acute liver injury that normally accompanies wild-type GBV-B or HCV infections. The demonstration of hepatitis in association with infection with the chimeric III^{HC} virus enhances its potential utility as a surrogate virus for evaluation of antiviral agents directed against the IRES or RNA replication signals residing within domain III of the HCV 5'NTR.

EXAMPLE 25

Sequence Analysis of Recovered Chimeric III^{HC} Virus

The lengthy delay prior to the appearance of high level viremia in T16444 coupled with the early robust replication of virus collected late in the course of infection in T16444 in a second animal suggests that the chimeric III^{HC} virus had undergone adaptation to the presence of the chimeric 5'NTR. To determine the nature of the mutations contributing to this adaptation, viral RNA from both animals was amplified by RT-PCR and subjected to sequencing and comparison with the parent synthetic III^{HC} RNA.

Viral RNA was isolated from serum obtained at week 14 from tamarin T16444, and week 4 or 8 from T16451 using Qiamp Viral RNA extraction kit (Qiagen). Random or GBV-B specific primers were used to prepare cDNA using MMLV reverse transcriptase from the Advantage RT-for-PCR kit (Clontech). PCR amplification of 1-2 kb fragments was carried out using the Advantage 2 PCR enzyme system (Clontech) utilizing GBV-B specific primers. Amplimers were purified through a silica gel membrane (QiaQuick PCR purification kit, Qiagen) and subjected to direct sequencing on an ABI 373XL instrument. The 3' terminus of the viral genome was ligated to a 27-mer oligonucleotide as described in the previous section for parent GBV-B. This RNA was then subjected to reverse transcription using a primer complementary to the ligated primer, and the cDNA was subsequently amplified using the same 3' primer and a 5' primer specific for a sequence in the 3'NTR (nts 9092-9111). A semi-nested PCR was performed with an internal 5' primer (nts 9097-9114). The 5' terminus of the viral genome was subjected to reverse transcription using a 3' primer complementary to the domain III of the HCV 5'NTR sequence (nts 209-229 of the HCV cDNA). The resulting cDNA was ligated to the 27-mer oligonucleotide and amplified using a primer complementary to the ligated primer and the same 3' primer used for RT. A semi-nested PCR was then performed using an internal 3' primer (nts 137-156 of the HCV cDNA). Results are shown in FIG. 7.

EXAMPLE 26**Use of the Chimeric IRES Virus GB/III^{HC} to Study the Efficacy of Potential Antiviral Candidates Targeting Domain III of the HCV IRES (III^{HC})**

GB/III^{HC} chimeric virus replicates robustly in small primates (see FIG. 11) and were
5 therefore used for antiviral validation studies in the small primate model. GB/III^{HC} chimeric virus were also demonstrated to replicate well in primary cultures of tamarin hepatocytes (FIG. 8).

Primary tamarin hepatocytes (*Saguinus mystax*) were prepared as described previously (Beames *et al.*, 2000). Wild-type (wt) GBV-B and chimeric virus GB/III^{HC} inocula were derived
10 from tamarins that had been infected with the corresponding viruses. 2.5×10^6 cells were seeded into each well of 6-well plates, and subsequently infected with virus (wt or chimera) at a multiplicity of infection (m.o.i.) of approximately 0.6 genome equivalents (ge)/cell. One hour after infection, cells were thoroughly washed 5 times and fresh medium was added. The media was replaced with fresh media every 24 hours thereafter. GBV-B RNA in culture supernatants
15 and cell extracts was measured quantitatively at the indicated times (FIG. 8) by a realtime RT-PCR (TaqMan®) assay targeting GBV-B core protein coding sequence (Beames *et al.*, 2000). Results showed that there is synthesis and secretion of newly formed viral particles in the culture supernatant after infection with both the wild-type and chimeric viruses. The chimeric IRES virus replicated efficiently in these cultures, although to titers that were about 10-fold less than
20 the wild-type GBV-B titers.

These data show that the GB/III^{HC} chimeric virus makes possible the validation of candidate antiviral compounds targeting domain III of the HCV IRES in the context of a complete viral replication cycle in primary hepatocytes in *ex vivo* culture. This is a significant
25 advantage over the only other system for cell culture validation of antiviral compounds, which consists of a cell culture system wherein only the effect on viral RNA replication can be monitored. The GB/III^{HC} chimeric virus also allows such studies in the small primate model: tamarins, marmosets, or owl monkeys.

EXAMPLE 27**Determination of the Mutations Contributing to Efficient Replication of the III^{HC} Chimeric Virus.**

Six mutations were identified in the viral RNA isolated at week 14 post-inoculation (p.i.)
30 from a tamarin (T16444) that was inoculated with the domain III chimeric RNA (III^{HC}) (FIG. 9). Two mutations were located in the 3'NTR, two were silent substitutions, in the E1 and NS5B

coding regions, respectively, and two were substitutions leading to amino acid changes in the E2 and NS5B sequences, respectively (FIG.10). The fact that high titers of virus appeared in the serum only many weeks after RNA inoculation of tamarin T16444, and that this virus contained these mutations, suggests that the mutations (either singly or collectively) may be required for robust replication of the chimeric virus.

Virus recovered from T16444, that contains the chimeric 5'NTR as well as all of these mutations, replicated efficiently when passaged in a second tamarin (T16451, FIG. 11).

To assess the mutations role in replication of III^{HC} chimera, these mutations were iteratively introduced into the cDNA of the initial III^{HC} chimera. Four plasmid DNAs were thus constructed (FIG. 12). After linearization of the plasmid DNAs by digestion with *Xho*I, genome-length RNA transcripts containing the chimeric 5'NTR and the indicated mutations elsewhere in the genome were transcribed from the cognate plasmids using the T7 MegaScript kit™ (Ambion). An aliquot of the reaction products was tested by agarose gel electrophoresis to ensure RNA integrity and to approximate the quantity of RNA. The remaining RNA was frozen at -80°C until inoculation, without further purification, into the liver of susceptible GBV-B naïve tamarins.

Individual animals (*S. mystax*) received a dose of approximately 100 µg of RNA transcribed from one of the plasmids shown in FIG. 12. The RNAs were inoculated into the liver under direct visualization following exposure by laparotomy. Inoculated animals were followed as described above, with periodic testing of serum ALT, antibody to the NS3 protein of GBV-B, and the presence of GBV-B RNA in serum. The tamarins were housed at the Southwest Regional Primate Research Center at the Southwest Foundation for Biomedical Research. They were cared for by members of the Department of Laboratory Animal Medicine of the Southwest Foundation for Biomedical Research in accordance with the Guide for the Care and Use of laboratory Animals. All protocols were approved by the Institutional Animal Care and Use Committee (IACUC).

The primary indicator of successful infection was detection of sustained viremia in sequential serum samples collected over a period of weeks. Viremia was monitored by detection of GBV-B RNA using a sensitive and specific real-time RT-PCR assay. RNA was isolated from virus present in tamarin serum using the QIAamp® viral RNA extraction kit (Qiagen) and amplified in a one-step RT-PCR reaction utilizing TaqMan® EZ core reagents (Perkin Elmer). First-strand cDNA synthesis was carried out using the primer NS5ARpp (5'-GAAGGAGGGAGGTTTGAAGGA-3' (SEQ ID NO:14), position 6949-6969). The cDNA

product was quantified in a 5' exonuclease PCR (TaqMan®) assay using a primer-probe combination that recognized sequence in the GBV-B NS5A coding region. The primers NS5ARPp and NS5AFPp (5'-CCAGTTCCGGGCAAGAA CT-3' (SEQ ID NO:15), position 6844-6862) and probe (nts 6913-6938) were obtained from Fisher and Perkin-Elmer, respectively. A synthetic RNA derived from the infectious clone was used as the standard for quantitation of the RNA.

RNAs containing only the NS5A mutation (GB/III^{HC}-m1) failed to replicate. On the other hand, in one of two inoculated animals, a chimeric RNA containing the NS5A mutation together with both 3'NTR mutations (GB/III^{HC}-m2) replicated in a robust fashion at week 4 post-inoculation, reaching viremia levels in excess of 10⁷ genome equivalents/ml (FIG. 12). Consistent with this latter result, the introduction of the E2 mutation into the mutated chimeric RNA containing the NS5A and 3'NTR mutations (GB/III^{HC}-m3), or a modified chimeric RNA containing all 6 mutations (GB/III^{HC}-m4) (FIG. 12), resulted in rapid rescue of virus and the presence of substantial viremia within 2 weeks of intrahepatic inoculation of the RNA (FIG. 13).

These data indicate that one or both of the two mutations in the 3' NTR sequence of GBV-B (C9052U and C9065U) are involved in an efficient-replication phenotype of the domain III chimeric virus. These mutations appear to be useful for efficient replication both *in vivo* in intrahepatically inoculated animals and *ex vivo* in cultured primary hepatocytes.

These data are strongly suggestive of a direct interaction between the 3'NTR in the region of nucleotides 9052 and 9065 and the 5'NTR in the region of domain III of the IRES. Such circularizing interactions of the termini of genomic RNA are well known to occur in other members of the *Flaviviridae* family and are essential for viral replication. However, they have not been previously documented to occur in hepaciviruses.

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents, which are both chemically and physiologically related, may

be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

5

U.S. Patent 4,578,770

U.S. Patent 4,596,792

U.S. Patent 4,599,230

U.S. Patent 4,599,231

10 U.S. Patent 4,601,903

U.S. Patent 4,608,251

U.S. Patent 4,683,195

U.S. Patent 4,683,202

U.S. Patent 4,800,159

15 U.S. Patent 4,883,750

U.S. Patent 5,279,721

U.S. Patent 5,807,670

EPA No. 320,308

EPA No. 329,822

20 GB Application No. 2,202,328

PCT/US87/00880

PCT/US89/01025

WO 88/10315

WO 89/06700

25 WO 90/07641

Al et al., Virus Res., 53:141-149, 1998.*Alter et al., Hepatology*, 26:625-655, 1997.*Beames et al., J. Virol.*, 74:11764-11772, 2000.*Beard et al., Hepatology*, 1999.

- Behrens *et al.*, *EMBO. J.*, 15:12-22, 1996.
- Chambers *et al.*, *J. Virol.*, 73:3095-3101, 1999.
- Choo *et al.*, *Proc. Natl. Acad. Sci. USA*, 91:1294-1298, 1994.
- Consensus Development Panel. National Institutes of Health Consensus Development
5 Conference Panel statement: Management of hepatitis C. *Hepatology*, 26:Supplement
1[3], 2S-10S, 1997.
- Deinhardt *et al.*, *J. Exp Med.* 125:637-688, 1967.
- Farci *et al.*, *Science*, 258:135-140. 1992.
- Farci *et al.*, *J. Infectious Diseases*, 165:1006-1011, 1992.
- 10 Filocamo *et al.*, *J. Virol.*, 71 1417-27, 1997.
- Frohman, *PCR Protocols: A Guide To Methods And Applications*, Academic Press, New York,
1990.
- Frolov *et al.*, *RNA*, 4:1418-1435, 1998.
- Grakoui *et al.*, *J. Virol.*, 67(5):2832-2843, 1993.
- 15 Grakoui *et al.*, *J. Virol.*, 67(3):1385-95. 1993.
- Hijikata *et al.*, *J. Virol.*, 67:4665-4675, 1993.
- Honda *et al.*, *RNA*, 2:955-956, 1996.
- Honda *et al.*, *J. Virol.*, 73:1165-1174, 1999.
- Inchauspe *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:10292-10296, 1991.
- 20 Karayiannis *et al.*, *Hepatology*, 9:186-192, 1989.
- Kim *et al.*, *Cell*, 87:343-355, 1996.
- Kolykhalov *et al.*, *Science*, 277:570-574, 1997.
- Kolykhalov *et al.*, *J. Virol.*, 70:3363-3371, 1996.
- Landt *et al.*, *Gene*, 96:125-128, 1990.
- 25 Lemon and Honda, *Seminars in Virology*, 8:274-288, 1997.
- Lindenbach and Rice, *J. Virol.*, 73:4611-4621, 1999.
- Lu and Wimmer, *Proc., Natl., Acad., Sci., USA*, 93:1412-1417, 1996.
- Muerhoff *et al.*, *J. Virol.*, 69:5621-5630, 1995.
- Neumann *et al.*, *Science*, 282:103-107, 1998.
- 30 Pletnev *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:10532-10536, 1992.
- Remington's Pharmaceutical Sciences, 15th Edition, pages 1035-1038 and 1570-1580.
- Reynolds *et al.*, *EMBO. J.*, 14:6010-6020, 1995.

- Rijnbrand *et al.*, Mutational analysis of the GB virus B internal ribosome entry site. *J. Virol.*, 74(2):773-783, 1999.
- Sambrook, *et al.*, *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring, Harbor, N.Y., 1989.
- 5 Scarselli *et al.*, *J. Virol.*, 71 4985-4989, 1997.
- Schlauder *et al.*, *J. Med. Virol.*, 46:81-90, 1995.
- Shaffer *et al.*, *J. Virol.*, 69:600-6604, 1995.
- Simons, *et al.*, *Proc. Natl. Acad. Sci.*, 92:3401-3405, 1995.
- Tanaka *et al.*, *Biochem. Biophys. Res. Comm.*, 215:744-749, 1995.
- 10 Teller *et al.*, *J. Clin. Micro.*, 34:3085-3091, 1996.
- Todd *et al.*, *J. Virol.*, 71:8868-74, 1997.
- Yanagi *et al.*, *Proc. Natl. Acad. Sci. USA*, 16:8738-8743, 1998.
- Yao *et al.*, *Nature Structural Biology*, 4: 463-467, 1997.
- Zhao *et al.*, *J. Virol.*, 73:1546-1554, 1999.

CLAIMS

1. An isolated polynucleotide comprising a chimeric GBV-B polynucleotide encoding a virus.
5
2. The polynucleotide of claim 1, comprising a chimeric GBV-B genome, wherein at least part, but not all of a 5' NTR sequence is from a HCV 5' NTR.
3. The polynucleotide of claim 2, wherein at least one, but not all of domain I, II, III, or IV
10 of the 5' NTR is from a HCV 5' NTR.
4. The polynucleotide of claim 2, wherein domain I of the 5' NTR is from a HCV 5'NTR.
5. The polynucleotide of claim 2, wherein domain II of the 5' NTR is from a HCV 5'NTR.
15
6. The polynucleotide of claim 2, wherein domain III of the 5' NTR is from a HCV 5'NTR.
7. The polynucleotide of claim 6, wherein the 5' NTR domain Ib of GBV-B is deleted.
- 20 8. The polynucleotide of claim 2, wherein domain IV of the 5' NTR is from a HCV 5'NTR.
9. The polynucleotide of claim 2, wherein domain I and domain II of the 5' NTR is from a HCV 5'NTR.
- 25 10. The polynucleotide of claim 2, wherein domain I and domain III of the 5' NTR is from a HCV 5'NTR.
11. The polynucleotide of claim 2, wherein domain I and domain IV of the 5' NTR is from a HCV 5'NTR.
30
12. The polynucleotide of claim 2, wherein domain II and domain III of the 5' NTR is from a HCV 5'NTR.

13. The polynucleotide of claim 2, wherein domain II and domain IV of the 5' NTR is from a HCV 5'NTR.
14. The polynucleotide of claim 2, wherein domain III and domain IV of the 5' NTR is from a HCV 5'NTR.
15. The polynucleotide of claim 2, wherein domain II, domain III and domain IV of the 5' NTR is from a HCV 5'NTR.
16. The polynucleotide of claim 15, wherein the 5' NTR domain Ib of GBV-B is deleted.
17. The polynucleotide of claim 2, wherein said polynucleotide is DNA.
18. The polynucleotide of claim 2, wherein said polynucleotide is RNA.
19. The polynucleotide of claim 1, further comprising at least part of a structural protein coding region of HCV.
20. The polynucleotide of claim 1, further comprising at least part of a non-structural protein coding region of HCV.
21. A viral expression construct comprising a chimeric GBV-B polynucleotide, wherein at least a part of the 5' NTR sequence is from a HCV 5' NTR.
22. The viral expression construct of claim 21, wherein domain III of the 5' NTR is from a HCV 5' NTR.
23. The viral expression construct of claim 21, further comprising a deletion of the GBV-B 5' NTR domain Ib region.
24. The viral expression construct of claim 21, wherein said construct is a plasmid.
25. The viral expression construct of claim 21, wherein said construct is a virus.

26. The viral expression construct of claim 21, further defined as a construct for the expression of a chimeric GBV-B/HCV virus.
27. A method for identifying a compound active against a viral infection comprising:
5 providing a virus expressed from a viral construct comprising a chimeric GBV-B/HCV virus;
contacting the virus with a candidate substance; and
comparing infectivity of the virus in the presence of the candidate substance with the infectivity of the virus in the absence of the candidate substance.
10
28. The method of claim 27, wherein the chimeric virus comprises at least part of a 5' NTR sequence from a HCV 5' NTR.
29. The method of claim 28, wherein the chimeric virus comprises domain III of the 5' NTR
15 is from a HCV 5'NTR.
30. The method of claim 28, wherein the chimeric virus comprises a deletion of domain Ib of GBV-B.
- 20 31. A compound active against a viral infection identified according to a method comprising:
providing a virus expressed from a viral construct comprising a GBV-B/HCV chimeric virus;
contacting the virus with a candidate substance; and
comparing the infectivity of the virus in the presence of said candidate substance with the
25 infectivity of the virus in the absence of said candidate substance.
32. The compound of claim 31, wherein the compound is identified by providing a viral construct comprising at least part of a 5' NTR sequence from a HCV 5' NTR.
- 30 33. The compound of claim 32, wherein the compound is identified by providing a viral construct comprising domain III of the 5' NTR is from a HCV 5'NTR.

34. The compound of claim 32, wherein the compound is identified by providing a viral construct lacks domain Ib of GBV-B.
35. A hepatotropic virus comprising a chimeric GBV-B polynucleotide, wherein at least a part of the 5' NTR sequence is from a HCV 5' NTR.
36. The hepatotropic virus of claim 35, wherein domain III of the 5' NTR is from a HCV 5' NTR.
37. The hepatotropic virus of claim 35, further comprising a deletion of the GBV-B 5' NTR domain Ib region.
38. The hepatotropic virus of claim 36, wherein the virus propagates *in vivo*.
39. The hepatotropic virus of claim 35, further defined as a construct for the expression of a chimeric GBV-B/HCV virus.
40. A method of producing a virus comprising:
introducing into a host cell a viral expression construct comprising a chimeric GBV-B polynucleotide encoding at least part of a 5' NTR sequence from a HCV 5' NTR sequence; and
culturing the host cell under conditions permitting production of a virus from the construct.
41. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain I from a HCV 5' NTR.
42. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain II from a HCV 5' NTR.
43. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain III from a HCV 5' NTR.

44. The method of claim 43, wherein said polynucleotide comprises a deletion of 5' NTR domain Ib of GBV-B.
45. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain I and domain II from a HCV 5' NTR.
46. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain I and domain IV from a HCV 5' NTR.
47. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain I, domain II and domain III from a HCV 5' NTR.
48. The method of claim 40, wherein said polynucleotide comprises a 5' NTR from a HCV 5' NTR.
49. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain II and domain III from a HCV 5' NTR.
50. The method of claim 40, wherein said polynucleotide comprises at least a 5' NTR domain III and domain IV from a HCV 5' NTR.
51. The method of claim 40, wherein said host cell is a eukaryotic cell.
52. The method of claim 51, wherein said host cell is in an animal.
53. The method of claim 40, wherein said polynucleotide comprises synthetic RNA.
54. The method of claim 40, further comprising the step of isolating virus from said host cell.
55. The method of claim 54, wherein the isolated virus is purified to homogeneity.
56. A method for identifying a compound active against a viral infection comprising:

providing a chimeric virus expressed from a viral construct comprising at least part of a
5' NTR from a HCV 5' NTR;
contacting the virus with a candidate substance; and
comparing the infectivity of the virus in the presence of the candidate substance with the
5 infectivity of the virus in the absence of the candidate substance.

57. The method of claim 56, wherein the viral construct comprises at least a 5' NTR domain
III from a HCV 5' NTR.

10 58. The method of claim 56, wherein the viral construct lacks a 5' NTR domain Ib of GBV-
B.

1/12




FIG. 1

2/12




FIG. 2

3/12




FIG. 4




FIG. 3

4/12




FIG. 5

5/12




FIG. 6

6/12

Sequence differences between chimeric viral RNAs recovered from animals T16444 and T16451, and the parental synthetic IIIHC RNA.

nt position	parent	T 16444	T 16451	protein / region	aa position	parent	T 16444	T 16451
		wk 14	wk 4				wk 14	wk 4
238**	C	-	Deleted	5'NTR	NA			
1012	G	A	A	E1	189	Glu	-	-
2376	T	C	C	E2	644	Ile	Thr	Thr
5225	G	-	A	NS4A	1594	Ala	-	Thr
6590	A	G / -	-	NS5A	2049	Lys	Glu / -	-
7152	T	C	C	NS5A	2236	Val	Ala	Ala
7945	A	G	G	NS5B	2500	Thr	-	-
8350	A	-	G	NS5B	2635	Thr	-	-
9052	C	T	T	3'NTR	NA			
9065	C	T	T	3'NTR	NA			
Note 1: Numbering according to GBV-B parent genome,								
except for ** change occurring at the 5' end of domain III of the HCV IRES								
Note 2: (-) identical to parent residue								

FIG. 7

7/12




FIG. 8




FIG. 9

8/12




FIG. 10




FIG. 11

9/12





FIG. 12

10/12






FIG. 14A

SEQUENCE LISTING

5 <110> MARTIN, ANNETTE
SANGAR, DAVID V.
LEMON, STANLEY M.
RIJNBRAND, RENE

10 <120> Chimeric GB Virus B (GBV-B)

<130> UTFG:258WO

<140> UNKNOWN

15 <141> 2003-07-02

<150> 10/189.359

<151> 2002-07-03

20 <160> 16

<170> PatentIn Ver. 2.1

<210> 1

25 <211> 260

<212> RNA

<213> Unknown

<220>

30 <223> Description of Unknown Organism: GB VIRUS B

<400> 1

gaguuuugcga ccaugggugga ucagaaccgu uucgggugaa gccauaggucu gaaggggaug 60

acgucuccuuc uggcucaucc acaaaaaccg ucucgggugg gugaggaguc cuggcugugu 120

35 gggaagcagu caguauaaau cccgucgugu guggugacgc cucacgacgu auuuguccgc 180

ugugcagagc guaguaccaa gggcugcacc ccgguuuuug uuccaagcgg agggcaaccc 240

ccgcuuggaa uaaaaaacug 260

40 <210> 2

<211> 9399

<212> DNA

<213> GBV-A-like virus

45 <400> 2

accacaaaca ctccagtttg ttacactccg ctagggaatgc tcctggagca cccccctag 60

cagggcgttg gggatttccc ctgcccgtct gcagaagggt ggagccaacc accttagtat 120

gtaggcggcg ggactcatga cgctcgctg atgacaagcg ccaagcttga cttggatggc 180

cctgatgggc gttcatgggt tcggtgggtg tggcgcttta ggcagcctcc acgcccacca 240

50 cctcccagat agagcggcgg cactgtaggg aagaccgggg accggtcact accaaggacg 300

cagacctctt tttgagtatc acgcctccgg aagtagttgg gcaagcccac ctatatgtgt 360

tgggatgggt ggggttagcc atccataccg tactgcctga tagggtcctt gcgaggggat 420

ctgggagtct cgtagaccgt agcacatgcc tgttatttct actcaaacia gtcctgtacc 480

55 tgcgccaga acgcgcaaga acaagcagac gcaggcttca tatcctgtgt ccattaaaac 540

atctgttgaa aggggacaac gagcaaagcg caaagtccag cgcgatgctc ggccctcgtaa 600

ttacaaaatt gctggtatcc atgatggctt gcagacattg gctcaggctg ctttgccagc 660

tcattggttg ggcagccaag accctcgcca taagtctcgc aatcttggaa tccttctgga 720

ttaccctttg ggggtgattg gtgatgttac aactcacaca cctctagtag gcccgtggtg 780

ggcaggagcg gtcgttcgac cagtctgcca gatagtacgc ttgctggagg atggagtcaa 840

60 ctgggctact gggttggttcg gtgtccacct ttttgtggta tgtctgctat ctttggcctg 900

tcctttagtg ggggcgcggg tctactgacc agacacaaat accacaatcc tgaccaattg 960

gtgccagcgt aatcagggtta tctattgttc tccttccact tgcctacacg agcctgggtg 1020

tgtgatctgt gcggacgagt gctgggttcc cgccaatccg tacatctcac acccttccaa 1080

65 ttggactggc acggactcct tcttggctga ccacattgat tttgttatgg gcgctcttgt 1140

gacctgtgac gcccttgaca ttggtgagtt gtgtgggtgcg tgtgtattag tcggtgactg 1200

gcttgtcagg cactggctta ttcacataga cctcaatgaa actggtactt gttacctgga 1260

	agtgccact	ggaatagatc	ctgggttccct	aggggtttatc	gggtggatgg	ccggcaaggt	1320
	cgaggctgtc	atcttcttga	ccaaactggc	ttcacaagta	ccatacgcta	ttgcgactat	1380
	gttttagcag	gtacactacc	tggcggttgg	cgctctgatc	tactatgcct	ctcggggcaa	1440
5	gtgggtatcag	ttgtctcctag	cgcttatgct	ttacatagaa	gcgacctctg	gaaaccctat	1500
	caggggtgccc	actggatgct	caatagctga	gttttgctcg	cctttgatga	taccatgtcc	1560
	ttgccactct	tattttgagt	agaatgtgtc	agaagtcat	tggtacagtc	caaagtggac	1620
	caggcctatc	actctagagt	ataacaactc	catatcttgg	tacccttata	caatccctgg	1680
	tgcgagggga	tgtatggtta	aattcaaaaa	taacacatgg	ggttgctgcc	gtattcgcaa	1740
10	tgtgccatcg	tactgcaacta	tgggcactga	tgcagtgtgg	aacgacactc	gcaacactta	1800
	cgaagtatgc	ggtgtaacac	catggctaac	aaccgcattg	cacaacggct	cagccctgaa	1860
	attggctata	ttacaatacc	ctgggtctaa	agaaatgttt	aaacctcata	attggatgtc	1920
	aggccatttg	tattttgagg	gatcagatac	ccctatagtt	tactttttatg	accctgtgaa	1980
	ttccactctc	ctaccaccgg	agaggtgggc	taggttgccc	ggtacccac	ctgtggtacg	2040
	tggttcttgg	ttacaggttc	cgcaagggtt	ttacagtgat	gtgaaagacc	tagccacagg	2100
15	attgatcacc	aaagacaaag	cctggaaaaa	ttatcagggtc	ttatatcccg	ccacgggtgc	2160
	tttgtctctt	acgggagtta	ccaccaaggc	cgtggtgcta	attctgttgg	ggttctgtgg	2220
	cagcaagtat	cttattttag	cctacctctg	cttctgttcc	ctttgttttg	ggcgcgcttc	2280
	tggttaccct	ttgcgtcctg	tgctcccatc	ccagtcgtat	ctccaagctg	gctgggatgt	2340
	tttgtctaaa	gctcaagtag	ctccttttgc	tttgattttc	ttcatctgtt	gctatctccg	2400
20	ctgcaggcta	cgttatgctg	cccttttagg	gtttgtgccc	atggctgcgg	gcttgccctt	2460
	aactttcttt	ggtgcagcag	ctgctgcccc	accagattat	gactggtggg	tgcgactgct	2520
	agtggcaggg	ttagttttgt	ggcgcgccg	taaccgtggt	caccgcatag	ctctgcttgt	2580
	aggtocttgg	cctctgttag	cgcttttaac	cctcttgcat	ttggttacgc	ctgcttcagc	2640
	ttttgatacc	gagataaattg	gagggctgac	aataccacct	gtagtagcat	tagttgtcat	2700
25	gtctcgtttt	ggcttctttg	ctcacttggt	acctcgctgt	gctttagtta	actcctatct	2760
	ttggcaacgt	tgggagaatt	ggttttggaa	cgttacacta	agaccggaga	ggtttttcct	2820
	tgtgctggtt	tgtttccccc	gtgcgacata	tgacgcgtg	gtgactttct	gtgtgtgtca	2880
	cgtagctctt	ctatgtttaa	catccagtgc	agcatcgctt	tttgggactg	actctagggt	2940
	tagggcccat	agaatgttgg	tgctctcgg	aaagtgtcat	gcttggtatt	ctcattatgt	3000
30	tcttaagttt	ttcctcttag	tgtttggtga	gaatggtgtg	tttttctata	agcacttgca	3060
	tgggtgatgtc	ttgcctaattg	attttgccctc	gaaactacca	ttgcaagagc	catttttccc	3120
	ttttgaaggc	aaggcaagg	tctataggaa	tgaaggaaga	cgcttgccgt	gtggggacac	3180
	ggttgatggt	ttgcccgttg	ttgcgcgtct	cggcgacctt	gttttcgcag	ggttggtctat	3240
	gcccgcagat	gggtgggcca	ttaccgcacc	ttttacgctg	cagtgtctct	ctgaacgtgg	3300
35	cacgctgtca	gcgatggcag	tggctcatgac	tggatatagc	ccccgaactt	ggactggaac	3360
	tatcttcaga	ttaggatctc	tggccactag	ctacatggga	tttgtttgtg	acaacgtgtt	3420
	gtatactgct	caccatggca	gcaaggggcg	ccggttggct	catcccacag	gctctataca	3480
	cccaataacc	ggtgacgcgg	ctaattgacca	ggacatctat	caaccaccat	gtggagctgg	3540
	gtcccttact	cggtgctctt	gcggggagac	caaggggtat	ctggtaacac	gactggggctc	3600
40	attggttgag	gtcaacaaat	ccgatgaccc	tatttgggtg	gtgtgcgggg	cccttcccat	3660
	ggctggtgcc	aagggttctt	caggtgcccc	gattctgtgc	tcctccgggc	atgttatttg	3720
	gatgttcacc	gctgctagaa	attctggcgg	ttcagtcagt	cagattagg	ttaggccgtt	3780
	ggtgtgtgct	ggataccatc	cccagtacac	agcacatgcc	actcttgata	caaaacctac	3840
	tgtgcctaac	gagtattcag	tgcaaatttt	aattgcccc	actggcagcg	gcaagtcaac	3900
45	caaattacca	ctttcttaca	tgcaggagaa	gtatgaggtc	ttggctctaa	atcccagttg	3960
	ggctacaaca	gcatcaatgc	caaagtacat	gcacgcgacg	tacggcgtga	atccaaattg	4020
	ctattttaat	ggcaaattgta	ccaacacagg	ggcttcaact	acgtacagca	catatggcat	4080
	gtacctgacc	ggagcatgtt	cccggaaacta	tgatgtaatc	atgtgtgacg	aatgccatgc	4140
	taccgatgca	accaccgtgt	tgggcatttg	aaaggctcta	accgaagctc	catccaaaaa	4200
50	tgttaggcta	gtggttcttg	ccacggctac	cccccttga	gtaatcccta	caccacatgc	4260
	caacataact	gagattcaat	taaccgatga	agggactatc	ccctttcatg	gaaaaaagat	4320
	taaggaggaa	aatctgaaga	aaggagagaca	ccctatcttt	gaggctacca	aaaaacactg	4380
	tgatgagctt	gctaaccgagt	tagctcgaaa	gggaataaca	gctgtctctt	actatagggg	4440
55	atgtgacatc	tcaaaaaatcc	ctgagggcga	ctgtgtagta	gttgccactg	atgccttgtg	4500
	tacagggtac	actggtgact	ttgattccgt	gtatgactgc	agcctcatgg	tagaaggcac	4560
	atgccattgt	gaccttgacc	ctacttttcac	ctaggtgtgt	cgtgtgtgctg	gggtttcagc	4620
	aatagttaaa	ggcgagcgta	ggggcgcgca	agggcgtggg	agagctggca	tataactata	4680
	tgtagacggg	agttgtaccc	cttcgggtat	ggttcctgaa	tgcaacattg	ttgaagcctt	4740
	cgacgcagcc	aaggcatggt	atggtttgtc	atcaacagaa	gctcaaaacta	ttctggacac	4800
60	ctatcgaccc	caacctgggt	tacctgcgat	aggagcaaat	ttggacgagt	gggctgatct	4860
	cttttctatg	gtcaaccccc	aaccttcatt	tgtcaatact	gcaaaaaagaa	ctgctgacaa	4920
	ttatgttttg	ttgactgcag	cccaactaca	actgtgtcat	cagtatggct	atgctgctcc	4980
	caatgacgca	ccacggtggc	agggagcccc	cttgggaaa	aaaccttgtg	gggttctgtg	5040
	gcgcttggac	ggcgctgacg	cctgtccttg	cccagagccc	agcgaggtga	ccagatacca	5100
65	aatgtgcttc	actgaagtca	atacttctgg	gacagccgca	ctcgctgttg	gcgttggagt	5160
	ggctatggct	tatctagcca	ttgacacttt	tggcgccact	tgtgtgcggc	gttgcgtgctc	5220

	tattgcatca	gtccctaccg	gtgctactgt	cgccccagtg	gttgacgaag	aagaaatcgt	5280
	ggaggagtgt	gcatcattca	ttcccttgga	ggccatgggt	gctgcaatcg	ataagctgaa	5340
	gagtacaatc	accacaacta	gtcctttcac	attggaaacc	gcccttgaaa	aacttaacac	5400
5	ctttcttggg	cctcatgcag	ctacaatcct	tgctatcata	gagtattgct	gtggtttagt	5460
	cactttacct	gacaatccct	ttgcatcatg	cgtgtttgct	ttcattgcgg	gtattactac	5520
	cccactacct	cacaagatca	aaatgttcct	gtcattatgt	ggaggcgcaa	ttgctgctcaa	5580
	gcttacagac	gctagaggcg	cactggcggt	catgatggcc	ggggctgcgg	gaacagctct	5640
	tgggtacatgg	acatcgggtg	gttttgtctt	tgacatgcta	ggcggctatg	ctgccgcctc	5700
	atccactgct	tgcttgacat	ttaaatgctt	gatgggtgag	tggccacta	tggatcagct	5760
10	tgtggtttt	gtctactccg	cgttcaatcc	ggcgcagga	gttgtggcg	tcttgtcagc	5820
	ttgtgcaatg	tttgctttga	caacagcagg	gccagatcac	tggcccaaca	gacttcttac	5880
	tatgcttgct	aggagcaaca	ctgtatgtaa	tgagtacttt	attgccactc	gtgacatccg	5940
	caggaagata	ctgggcatte	tggaggcatc	taccccttgg	agtgtcatat	cagcttgcat	6000
	ccgttggtct	cacaccccga	cggaggatga	ttgcggcctc	attgcttggg	gtctagagat	6060
15	ttggcagtat	gtgtgcaatt	tctttgtgat	ttgctttaat	gtccttaaag	ctggagttca	6120
	gagcatgggt	aacattcctg	gttgctcctt	ctacagctgc	cagaaggggt	acaagggccc	6180
	ctggatttga	tcaggatatgc	tccaagcacg	ctgtccatgc	gggtgctgaac	tcactctttc	6240
	tgttgagaat	ggttttgcaa	aactttacaa	aggaccaga	acttggtcaa	attactggag	6300
	aggggctgtt	ccagtcaacg	ctaggctgtg	tgggtcggct	agaccggacc	caactgattg	6360
20	gactagtctt	gtcgtcaatt	atggcggttag	ggactactgt	aaatatgaga	aaatgggaga	6420
	tcacattttt	gttacagcag	tatcctctcc	aaatgtctgt	ttcaccacag	tgcccccaac	6480
	cttgagagct	gcagtggcgg	tggacggcgt	acaggttcag	tgttatctag	gtgagcccaa	6540
	aactccttgg	acgacatctg	cttgctgtta	cggtcctgac	ggtaagggta	aaactgttaa	6600
	gcttcccttc	cgcgttgacg	gtcacacacc	tgggtgtgcg	atgcaactta	atttgctgta	6660
25	tgcacttgag	acaaatgact	gtaattccac	aaacaacact	cctagtgatg	aagccgcagt	6720
	gtccgctctt	gttttcaaac	aggagtgtcg	gcgtacaaac	caattgcttg	aggcaatttc	6780
	agctggcggt	gacaccacca	aactggcagc	ccccctccat	gaagaggtag	tggtaagaaa	6840
	gcgccagttc	cgggcaagaa	ctggttcgct	taccttgcc	ccccctccga	gatccgtccc	6900
	aggagtgtca	tgtcctgaaa	gcctgcaacg	aagtgaccgg	ttagaagggt	cttcaaacct	6960
30	ccctccttca	ccacctgttc	tacagttggc	catgccgatg	cccctgttgg	gagcgggtga	7020
	gtgtaaccct	ttcactgcaa	ttggatgtgc	aatgaccgaa	acaggcggag	gccctgatga	7080
	tttaccaggt	taccctccca	aaaaggaggt	ctctgaatgg	tcagacgaaa	gttggctcgac	7140
	ggctacaacc	gtttccagct	acgttactgg	ccccccgtac	cctaagatac	ggggaaagga	7200
	ttccactcag	tcagcccccg	ccaaacggcc	tacaaaaaag	aagttgggaa	agagtgaatt	7260
35	ttcgtgcagc	atgagctaca	cctggaccga	cgtgattagc	ttcaaaactg	cttctaaagt	7320
	tctgtctgca	actcggggcca	tcactagtgg	tttctcctca	caaagatcat	tgggtgatgt	7380
	gactgagccg	cgggatgcgg	agcttagaaa	acaaaaagtc	actattaata	gacaacctct	7440
	gttcccccca	tcataaccaca	agcaagttag	attggctaag	gaaaaagctt	caaaagttgt	7500
	cgggtgtcatg	tgggactatg	atgaagttag	agctcacacg	ccctctaagt	ctgctaagtc	7560
40	ccacatcaat	ggccttcggg	gcactgatgt	tcgttctgga	gcagcccgcg	aggctgttct	7620
	ggacttgacg	aagtgtgtcg	aggcagggtga	gataccgagt	cattatcggc	aaactgtgat	7680
	agttccaaag	gaggaggtct	tcgtgaagac	ccccagaaa	ccaacaaaga	aacccccaa	7740
	gcttatctcg	tacccccacc	ttgaaatgag	atgtgttgag	aagatgtact	acggtcaggt	7800
	tgtcctctgac	gtagttaaag	ctgtcatggg	agatgcgtac	gggtttgtag	atccacgtac	7860
45	ccgtgtcaag	cgtctgttgt	cgatgtggtc	acccgatgca	gtcggagcca	catgcgatac	7920
	agtggtgttt	gacagtagca	tcacaccgga	ggatatcatg	gtggagacag	acatctactc	7980
	agcagctaaa	ctcagtgacc	aacaccgagc	tggcattcac	accattgcga	ggcagttata	8040
	cgctggagga	ccgatgatcg	cttatgatgg	ccgagagatc	ggatatcgta	ggtgtagggt	8100
	ttccggcgct	tatactacct	caagttccaa	cagtttgacc	tgtctggctga	aggtaaatgc	8160
50	tgcagccgaa	caggctggca	tgaagaacct	tcgcttccct	atttgcggcg	atgattgcac	8220
	cgtaatttgg	aagagcgccg	gagcagatgc	agacaaacaa	gcaatgcgtg	tctttgctag	8280
	ctggatgaag	gtgatgggtg	caccacaaga	ttgtgtgcct	caacccaaat	acagttttga	8340
	agaattaaca	tcatgtctcat	caaagtgttac	ctctggaatt	accaaagtg	gcaagcctta	8400
	ctactttctt	acaagagatc	ctcgtatccc	ccttggcagg	tgtctgtccg	agggctctgg	8460
55	atacaacccc	agtgtctcgt	ggattgggtga	tctaatacat	cactacccat	gtttgtgggt	8520
	tagccgtgtg	tttgctgtcc	atttcatgga	gcagatgctc	tttgaggaca	aacttcccga	8580
	gacggtgacc	tttgactggg	atgggaaaaa	ttatagcgtg	cctgtagaag	atctgcccag	8640
	catcattgct	gggtgtgcacg	gtattgaggc	tttctcgggt	gtgcgctaca	ccaacgctga	8700
	gatcctcaga	gtttcccaat	cactaacaga	catgaccatg	ccccccctgc	gagcctggcg	8760
60	aaagaaagcc	agggcggtcc	tcgccagcgc	caagaggcgt	ggcggagcac	acgcaaaatt	8820
	ggctcgcttc	cttctctggc	atgctacatc	tagacctcta	ccagatttgg	ataagacgag	8880
	cgtggctcgg	tacaccactt	tcaattattg	tgtgttttac	tccccggagg	gggatgtgtt	8940
	tattacacca	cagagaagat	tgcagaagtt	tcttgtgaag	tatttggctg	tcattgtttt	9000
	tgccttaggg	ctcattgctg	ttggattagc	catcagctga	acccccaaat	tcaaaattaa	9060
65	ctaacagttt	tttttttttt	tttttttttt	agggcagcgg	caacagggga	gaccccgggc	9120
	ttaacgaccc	cgcgatgtg	agtttggcga	ccatgggtgga	tcagaacctg	ttcgggtgaa	9180

gccatggtct gaaggggatg acgtcccttc tggctcatcc acaaaaaccg tctcgggtgg 9240
gtgaggagtc ctggctgtgt gggagcagc cagtataatt cccgtcgtgt gtggtgacgc 9300
ctcacgacgt atttgtccgc tgtgcagagc gtagtaccaa gggctgcacc ccggtttttg 9360
ttccaagcgg agggcaaccc ccgcttgga ttaaaaact 9399

5

<210> 3

<211> 48

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

15

<400> 3

cgggatcccg taatacgact cactatagac cacaaacact ccagtttg

48

<210> 4

20 <211> 20

<212> DNA

<213> Artificial Sequence

<220>

25 <223> Description of Artificial Sequence: Primer

<400> 4

gtggaattca cagcgtcata

20

30

<210> 5

<211> 21

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence: Primer

<400> 5

40

tgtgaattcc actctcctac c

21

<210> 6

<211> 21

45 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

50

<400> 6

ttatcgattg cagcaaccat g

21

<210> 7

<211> 39

<212> DNA

<213> Artificial Sequence

60

<220>

<223> Description of Artificial Sequence: Primer

<400> 7

65

catggttgct gcaatcgata agctgaagag tacaataac

39

5 <210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

10 <400> 8
gacaacagac gcttgacacg 20

15 <210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

20 <400> 9
ctgtcatggg agatgcgtac 20

25 <210> 10
<211> 42
<212> DNA
<213> Artificial Sequence

30 <220>
<223> Description of Artificial Sequence: Primer

35 <400> 10
cgagctcgag cacatcgcg ggtcggttaag cccggggtct cc 42

40 <210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

45 <400> 11
gacaacagac gcttgacacg 20

50 <210> 12
<211> 108
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

55 <400> 12
ccgactcgag aattcggccc tgcaggccac aacagtctcg cgagttttta attccaagcg 60
ggggttgccc tccgcttgga acaaaaaccg ggggtgcagcc cttggtac 108

60

65 <210> 13
<211> 9599
<212> DNA
<213> Hepatitis C virus

<220>

<221> CDS

<222> (342) .. (9377)

5 <400> 13
gccagccccc tgatgggggc gacactccac catgaatcac tcccctgtga ggaactactg 60
tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgcgtgcag cctccaggac 120
10 cccccctccc gggagagcca tagtgggtctg cggaaccggt gactacaccg gaattgccag 180
gacgaccggg tcttttcttg gataaaccgg ctcaatgcct ggagatttgg gcgtgcccc 240
gcaagactgc tagccgagta gtgttgggtc gcgaaaggcc ttgtgggtact gcctgatagg 300
15 gtgcttgcca gtgccccggg aggtctcgta gaccgtgcac c atg agc acg aat cct 356
Met Ser Thr Asn Pro
1 5

20 aaa cct caa aga aaa acc aaa cgt aac acc aac cgt cgc cca cag gac 404
Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn Arg Arg Pro Gln Asp
10 15 20

25 gtc aag ttc ccg ggt ggc ggt cag atc gtt ggt gga gtt tac ttg ttg 452
Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly Gly Val Tyr Leu Leu
25 30 35

30 ccg cgc agg ggc cct aga ttg ggt gtg cgc gcg acg agg aag act tcc 500
Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala Thr Arg Lys Thr Ser
40 45 50

35 gag cgg tcg caa cct cga ggt aga cgt cag cct atc ccc aag gca cgt 548
Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro Ile Pro Lys Ala Arg
55 60 65

40 cgg ccc gag ggc agg acc tgg gct cag ccc ggg tac cct tgg ccc ctc 596
Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly Tyr Pro Trp Pro Leu
70 75 80 85

45 tat ggc aat gag ggt tgc ggg tgg gcg gga tgg ctc ctg tct ccc cgt 644
Tyr Gly Asn Glu Gly Cys Gly Trp Ala Gly Trp Leu Leu Ser Pro Arg
90 95 100

50 ggc tct cgg cct agc tgg ggc ccc aca gac ccc cgg cgt agg tcg cgc 692
Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro Arg Arg Arg Ser Arg
105 110 115

55 aat ttg ggt aag gtc atc gat acc ctt acg tgc ggc ttc gcc gac ctc 740
Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
120 125 130

60 atg ggg tac ata ccg ctc gtc ggc gcc cct ctt gga ggc gct gcc agg 788
Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
135 140 145

65 gcc ctg gcg cat ggc gtc cgg gtt ctg gaa gac ggc gtg aac tat gca 836
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala
150 155 160 165

70 aca ggg aac ctt cct ggt tgc tct ttc tct atc ttc ctt ctg gcc ctg 884
Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu
170 175 180

75 ctc tct tgc ctg act gtg ccc gct tca gcc tac caa gtg cgc aat tcc 932
Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Gln Val Arg Asn Ser
185 190 195

5	tcg ggg ctt tac cat gtc acc aat gat tgc cct aac tcg agt att gtg 980 Ser Gly Leu Tyr His Val Thr Asn Asp Cys Pro Asn Ser Ser Ile Val 200 205 210	
	tac gag gcg gcc gat gcc atc ctg cac act ccg ggg tgt gtc cct tgc 1028 Tyr Glu Ala Ala Asp Ala Ile Leu His Thr Pro Gly Cys Val Pro Cys 215 220 225	
10	gtt cgc gag ggt aac gcc tcg agg tgt tgg gtg gcg gtg acc ccc acg 1076 Val Arg Glu Gly Asn Ala Ser Arg Cys Trp Val Ala Val Thr Pro Thr 230 235 240 245	
15	gtg gcc acc agg gac ggc aaa ctc ccc aca acg cag ctt cga cgt cat 1124 Val Ala Thr Arg Asp Gly Lys Leu Pro Thr Thr Gln Leu Arg Arg His 250 255 260	
20	atc gat ctg ctt gtc ggg agc gcc acc ctc tgc tcg gcc ctc tac gtg 1172 Ile Asp Leu Leu Val Gly Ser Ala Thr Leu Cys Ser Ala Leu Tyr Val 265 270 275	
25	ggg gac ctg tgc ggg tct gtc ttt ctt gtt ggt caa ctg ttt acc ttc 1220 Gly Asp Leu Cys Gly Ser Val Phe Leu Val Gly Gln Leu Phe Thr Phe 280 285 290	
30	tct ccc agg cgc cac tgg acg acg caa gac tgc aat tgt tct atc tat 1268 Ser Pro Arg Arg His Trp Thr Thr Gln Asp Cys Asn Cys Ser Ile Tyr 295 300 305	
35	ccc ggc cat ata acg ggt cat cgc atg gca tgg gat atg atg atg aac 1316 Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn 310 315 320 325	
40	tgg tcc cct acg gca gcg ttg gtg gta gct cag ctg ctc cgg atc cca 1364 Trp Ser Pro Thr Ala Ala Leu Val Val Ala Gln Leu Leu Arg Ile Pro 330 335 340	
45	caa gcc atc atg gac atg atc gct ggt gct cac tgg gga gtc ctg gcg 1412 Gln Ala Ile Met Asp Met Ile Ala Gly Ala His Trp Gly Val Leu Ala 345 350 355	
50	ggc ata gcg tat ttc tcc atg gtg ggg aac tgg gcg aag gtc ctg gta 1460 Gly Ile Ala Tyr Phe Ser Met Val Gly Asn Trp Ala Lys Val Leu Val 360 365 370	
55	gtg ctg ctg cta ttt gcc ggc gtc gac gcg gaa acc cac gtc acc ggg 1508 Val Leu Leu Leu Phe Ala Gly Val Asp Ala Glu Thr His Val Thr Gly 375 380 385	
60	gga aat gcc ggc cgc acc acg gct ggg ctt gtt ggt ctc ctt aca cca 1556 Gly Asn Ala Gly Arg Thr Thr Ala Gly Leu Val Gly Leu Leu Thr Pro 390 395 400 405	
65	ggc gcc aag cag aac atc caa ctg atc aac acc aac ggc agt tgg cac 1604 Gly Ala Lys Gln Asn Ile Gln Leu Ile Asn Thr Asn Gly Ser Trp His 410 415 420	
70	atc aat agc acg gcc ttg aat tgc aat gaa agc ctt aac acc ggc tgg 1652 Ile Asn Ser Thr Ala Leu Asn Cys Asn Glu Ser Leu Asn Thr Gly Trp 425 430 435	
75	tta gca ggg ctc ttc tat caa cac aaa ttc aac tct tca ggc tgt cct 1700 Leu Ala Gly Leu Phe Tyr Gln His Lys Phe Asn Ser Ser Gly Cys Pro 440 445 450	
80	gag agg ttg gcc agc tgc cga cgc ctt acc gat ttt gcc cag ggc tgg 1748	

	Glu	Arg	Leu	Ala	Ser	Cys	Arg	Arg	Leu	Thr	Asp	Phe	Ala	Gln	Gly	Trp	
	455						460					465					
5	ggc	cct	atc	agt	tat	gcc	aac	gga	agc	ggc	ctc	gac	gaa	cgc	ccc	tac	1796
	Gly	Pro	Ile	Ser	Tyr	Ala	Asn	Gly	Ser	Gly	Leu	Asp	Glu	Arg	Pro	Tyr	
	470					475					480					485	
10	tgc	tgg	cac	tac	cct	cca	aga	cct	tgt	ggc	att	gtg	ccc	gca	aag	agc	1844
	Cys	Trp	His	Tyr	Pro	Pro	Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Lys	Ser	
					490					495					500		
15	gtg	tgt	ggc	ccg	gta	tat	tgc	ttc	act	ccc	agc	ccc	gtg	gtg	gtg	gga	1892
	Val	Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	
				505				510					515				
20	acg	acc	gac	agg	tcg	ggc	gcg	cct	acc	tac	agc	tgg	ggc	gca	aat	gat	1940
	Thr	Thr	Asp	Arg	Ser	Gly	Ala	Pro	Thr	Tyr	Ser	Trp	Gly	Ala	Asn	Asp	
			520					525					530				
25	acg	gat	gtc	ttc	gtc	ctt	aac	aac	acc	agg	cca	ccg	ctg	ggc	aat	tgg	1988
	Thr	Asp	Val	Phe	Val	Leu	Asn	Asn	Thr	Arg	Pro	Pro	Leu	Gly	Asn	Trp	
		535					540					545					
30	ttc	ggc	tgt	acc	tgg	atg	aac	tca	act	gga	ttc	acc	aaa	gtg	tgc	gga	2036
	Phe	Gly	Cys	Thr	Trp	Met	Asn	Ser	Thr	Gly	Phe	Thr	Lys	Val	Cys	Gly	
	550					555					560					565	
35	gcg	ccc	cct	tgt	gtc	atc	gga	ggg	gtg	ggc	aac	aac	acc	ttg	ctc	tgc	2084
	Ala	Pro	Pro	Cys	Val	Ile	Gly	Gly	Val	Gly	Asn	Asn	Thr	Leu	Leu	Cys	
					570					575					580		
40	ccc	act	gat	tgc	ttc	cgc	aaa	cat	ccg	gaa	gcc	aca	tac	tct	cgg	tgc	2132
	Pro	Thr	Asp	Cys	Phe	Arg	Lys	His	Pro	Glu	Ala	Thr	Tyr	Ser	Arg	Cys	
				585					590					595			
45	ggc	tcc	ggc	ccc	tgg	att	aca	ccc	agg	tgc	atg	gtc	gac	tac	ccg	tat	2180
	Gly	Ser	Gly	Pro	Trp	Ile	Thr	Pro	Arg	Cys	Met	Val	Asp	Tyr	Pro	Tyr	
			600					605				610					
50	agg	ctt	tgg	cac	tat	cct	tgt	acc	atc	aat	tac	acc	ata	ttc	aaa	gtc	2228
	Arg	Leu	Trp	His	Tyr	Pro	Cys	Thr	Ile	Asn	Tyr	Thr	Ile	Phe	Lys	Val	
		615					620					625					
55	agg	atg	tac	gtg	gga	ggg	gtc	gag	cac	agg	ctg	gaa	gag	gac	agg	tgc	2276
	Arg	Met	Tyr	Val	Gly	Gly	Val	Glu	His	Arg	Leu	Glu	Ala	Ala	Cys	Asn	
	630					635					640					645	
60	tgg	acg	cgg	ggc	gaa	cgc	tgt	gat	ctg	gaa	gac	agg	gac	agg	tcc	gag	2324
	Trp	Thr	Arg	Gly	Glu	Arg	Cys	Asp	Leu	Glu	Asp	Arg	Asp	Arg	Ser	Glu	
					650					655					660		
65	ctc	agc	ccg	ttg	ctg	ctg	tcc	acc	aca	cag	tgg	cag	gtc	ctt	ccg	tgt	2372
	Leu	Ser	Pro	Leu	Leu	Leu	Ser	Thr	Thr	Gln	Trp	Gln	Val	Leu	Pro	Cys	
				665				670					675				
70	tct	ttc	acg	acc	ctg	cca	gcc	ttg	tcc	acc	ggc	ctc	atc	cac	ctc	cac	2420
	Ser	Phe	Thr	Thr	Leu	Pro	Ala	Leu	Ser	Thr	Gly	Leu	Ile	His	Leu	His	
			680					685					690				
75	cag	aac	att	gtg	gac	gtg	cag	tac	ttg	tac	ggg	gta	ggg	tca	agc	atc	2468
	Gln	Asn	Ile	Val	Asp	Val	Gln	Tyr	Leu	Tyr	Gly	Val	Gly	Ser	Ser	Ile	
		695					700					705					
80	gcg	tcc	tgg	gcc	att	aag	tgg	gag	tac	gtc	gtt	ctc	ctg	ttc	ctt	ctg	2516
	Ala	Ser	Trp	Ala	Ile	Lys	Trp	Glu	Tyr	Val	Val	Leu	Leu	Phe	Leu	Leu	
	710					715					720					725	

5	ctt gca gac gcg cgc gtc tgc tcc tgc ttg tgg atg atg tta ctc ata	2564
	Leu Ala Asp Ala Arg Val Cys Ser Cys Leu Trp Met Met Leu Leu Ile	730 735 740
10	tcc caa gcg gag gcg gct ttg gag aac ctc gta ata ctc aat gca gca	2612
	Ser Gln Ala Glu Ala Ala Leu Glu Asn Leu Val Ile Leu Asn Ala Ala	745 750 755
15	tcc ctg gcc ggg acg cac ggt ctt gtg tcc ttc ctc gtg ttc ttc tgc	2660
	Ser Leu Ala Gly Thr His Gly Leu Val Ser Phe Leu Val Phe Phe Cys	760 765 770
20	ttt gcg tgg tat ctg aag ggt agg tgg gtg ccc gga gcg gtc tac gcc	2708
	Phe Ala Trp Tyr Leu Lys Gly Arg Trp Val Pro Gly Ala Val Tyr Ala	775 780 785
25	ctc tac ggg atg tgg cct ctc ctc ctg ctc ctg ctg gcg ttg cct cag	2756
	Leu Tyr Gly Met Trp Pro Leu Leu Leu Leu Leu Leu Ala Leu Pro Gln	790 795 800 805
30	cgg gca tac gca ctg gac acg gag gtg gcc gcg tcg tgt ggc ggc gtt	2804
	Arg Ala Tyr Ala Leu Asp Thr Glu Val Ala Ala Ser Cys Gly Gly Val	810 815 820
35	gtt ctt gtc ggg tta atg gcg ctg act ctg tcg cca tat tac aag cgc	2852
	Val Leu Val Gly Leu Met Ala Leu Thr Leu Ser Pro Tyr Tyr Lys Arg	825 830 835
40	tat atc agc tgg tgc atg tgg tgg ctt cag tat ttt ctg acc aga gta	2900
	Tyr Ile Ser Trp Cys Met Trp Trp Leu Gln Tyr Phe Leu Thr Arg Val	840 845 850
45	gaa gcg caa ctg cac gtg tgg gtt ccc ccc ctc aac gtc cgg ggg ggg	2948
	Glu Ala Gln Leu His Val Trp Val Pro Pro Leu Asn Val Arg Gly Gly	855 860 865
50	cgc gat gcc gtc atc tta ctc atg tgt gta gta cac ccg acc ctg gta	2996
	Arg Asp Ala Val Ile Leu Leu Met Cys Val Val His Pro Thr Leu Val	870 875 880 885
55	ttt gac atc acc aaa cta ctc ctg gcc atc ttc gga ccg ctt tgg att	3044
	Phe Asp Ile Thr Lys Leu Leu Leu Ala Ile Phe Gly Pro Leu Trp Ile	890 895 900
60	ctt caa gcc agt ttg ctt aaa gtc ccc tac ttc gtg cgc gtt caa ggc	3092
	Leu Gln Ala Ser Leu Leu Lys Val Pro Tyr Phe Val Arg Val Gln Gly	905 910 915
65	ctt ctc cgg atc tgc gcg cta gcg cgg aag ata gcc gga ggt cat tac	3140
	Leu Leu Arg Ile Cys Ala Leu Ala Arg Lys Ile Ala Gly Gly His Tyr	920 925 930
70	gtg caa atg gcc atc atc aag tta ggg gcg ctt act ggc acc tat gtg	3188
	Val Gln Met Ala Ile Ile Lys Leu Gly Ala Leu Thr Thr Tyr Val	935 940 945
75	tat aac cat ctc acc cct ctt cga gac tgg gcg cac aac ggc ctg cga	3236
	Tyr Asn His Leu Thr Pro Leu Arg Asp Trp Ala His Asn Gly Leu Arg	950 955 960 965
80	gat ctg gcc gtg gct gtg gaa cca gtc gtc ttc tcc cga atg gag acc	3284
	Asp Leu Ala Val Ala Val Glu Pro Val Val Phe Ser Arg Met Glu Thr	970 975 980
85	aag ctc atc acg tgg ggg gca gat acc gcc gcg tgc ggt gac atc atc	3332

	Lys	Leu	Ile	Thr	Trp	Gly	Ala	Asp	Thr	Ala	Ala	Cys	Gly	Asp	Ile	Ile	
				985					990					995			
5	aac	ggc	ttg	ccc	gtc	tct	gcc	cgt	agg	ggc	cag	gag	ata	ctg	ctt	ggg	3380
	Asn	Gly	Leu	Pro	Val	Ser	Ala	Arg	Arg	Gly	Gln	Glu	Ile	Leu	Leu	Gly	
		1000						1005					1010				
10	cca	gcc	gac	gga	atg	gtc	tcc	aag	ggg	tgg	agg	ttg	ctg	gcg	ccc	atc	3428
	Pro	Ala	Asp	Gly	Met	Val	Ser	Lys	Gly	Trp	Arg	Leu	Leu	Ala	Pro	Ile	
		1015					1020					1025					
15	acg	gcg	tac	gcc	cag	cag	acg	aga	ggc	ctc	cta	ggg	tgt	ata	atc	acc	3476
	Thr	Ala	Tyr	Ala	Gln	Gln	Thr	Arg	Gly	Leu	Leu	Gly	Cys	Ile	Ile	Thr	
	1030					1035					1040					1045	
20	agc	ctg	act	ggc	cgg	gac	aaa	aac	caa	gtg	gag	ggg	gag	gtc	cag	atc	3524
	Ser	Leu	Thr	Gly	Arg	Asp	Lys	Asn	Gln	Val	Glu	Gly	Glu	Val	Gln	Ile	
				1050					1055						1060		
25	gtg	tca	act	gct	acc	caa	acc	ttc	ctg	gca	acg	tgc	atc	aat	ggg	gta	3572
	Val	Ser	Thr	Ala	Thr	Gln	Thr	Phe	Leu	Ala	Thr	Cys	Ile	Asn	Gly	Val	
				1065					1070						1075		
30	tgc	tgg	act	gtc	tac	cac	ggg	gcc	gga	acg	agg	acc	atc	gca	tca	ccc	3620
	Cys	Trp	Thr	Val	Tyr	His	Gly	Ala	Gly	Thr	Arg	Thr	Ile	Ala	Ser	Pro	
		1080						1085					1090				
35	aag	ggg	cct	gtc	atc	cag	atg	tat	acc	aat	gtg	gac	caa	gac	ctt	gtg	3668
	Lys	Gly	Pro	Val	Ile	Gln	Met	Tyr	Thr	Asn	Val	Asp	Gln	Asp	Leu	Val	
		1095					1100					1105					
40	ggc	tgg	ccc	gct	cct	caa	ggg	tcc	cgc	tca	ttg	aca	ccc	tgt	acc	tgc	3716
	Gly	Trp	Pro	Ala	Pro	Gln	Gly	Ser	Arg	Ser	Leu	Thr	Pro	Cys	Thr	Cys	
	1110					1115					1120					1125	
45	ggc	tcc	tcg	gac	ctt	tac	ctg	gtc	acg	agg	cac	gcc	gat	gtc	att	ccc	3764
	Gly	Ser	Ser	Asp	Leu	Tyr	Leu	Val	Thr	Arg	His	Ala	Asp	Val	Ile	Pro	
				1130						1135					1140		
50	gtg	cgc	cgg	cga	ggg	gat	agc	agg	ggg	agc	ctg	ctt	tcg	ccc	cgg	ccc	3812
	Val	Arg	Arg	Arg	Gly	Asp	Ser	Arg	Gly	Ser	Leu	Leu	Ser	Pro	Arg	Pro	
				1145				1150						1155			
55	att	tcc	tac	ttg	aaa	ggc	tcc	tcg	ggg	ggg	ccg	ctg	ttg	tgc	ccc	gcg	3860
	Ile	Ser	Tyr	Leu	Lys	Gly	Ser	Ser	Gly	Gly	Pro	Leu	Leu	Cys	Pro	Ala	
			1160					1165						1170			
60	gga	cac	gcc	gtg	ggc	cta	ttc	agg	gcc	gcg	gtg	tgc	acc	cgt	gga	gtg	3908
	Gly	His	Ala	Val	Gly	Leu	Phe	Arg	Ala	Ala	Val	Cys	Thr	Arg	Gly	Val	
		1175					1180					1185					
65	gct	aaa	gcg	gtg	gac	ttt	atc	cct	gtg	gag	aac	cta	ggg	aca	acc	atg	3956
	Ala	Lys	Ala	Val	Asp	Phe	Ile	Pro	Val	Glu	Asn	Leu	Gly	Thr	Thr	Met	
	1190					1195					1200					1205	
70	aga	tcc	ccg	gtg	ttc	acg	gac	aac	tcc	tct	cca	cca	gca	gtg	ccc	cag	4004
	Arg	Ser	Pro	Val	Phe	Thr	Asp	Asn	Ser	Ser	Pro	Pro	Ala	Val	Pro	Gln	
				1210						1215					1220		
75	agc	ttc	cag	gtg	gcc	cac	ctg	cat	gct	ccc	acc	ggc	agc	ggg	aag	agc	4052
	Ser	Phe	Gln	Val	Ala	His	Leu	His	Ala	Pro	Thr	Gly	Ser	Gly	Lys	Ser	
				1225					1230						1235		
80	acc	aag	gtc	ccg	gct	gcg	tac	gca	gcc	cag	ggc	tac	aag	gtg	ttg	gtg	4100
	Thr	Lys	Val	Pro	Ala	Ala	Tyr	Ala	Ala	Gln	Gly	Tyr	Lys	Val	Leu	Val	
			1240					1245						1250			

	ctc aac ccc tct gtt gct gca acg ctg ggc ttt ggt gct tac atg tcc	4148
	Leu Asn Pro Ser Val Ala Ala Thr Leu Gly Phe Gly Ala Tyr Met Ser	
	1255 1260 1265	
5	aag gcc cat ggg gtt gat cct aat atc agg acc ggg gtg aga aca att	4196
	Lys Ala His Gly Val Asp Pro Asn Ile Arg Thr Gly Val Arg Thr Ile	
	1270 1275 1280 1285	
10	acc act ggc agc ccc atc acg tac tcc acc tac ggc aag ttc ctt gcc	4244
	Thr Thr Gly Ser Pro Ile Thr Tyr Ser Thr Tyr Gly Lys Phe Leu Ala	
	1290 1295 1300	
15	gac ggc ggg tgc tca gga ggt gct tat gac ata ata att tgt gac gag	4292
	Asp Gly Gly Cys Ser Gly Gly Ala Tyr Asp Ile Ile Ile Cys Asp Glu	
	1305 1310 1315	
20	tgc cac tcc acg gat gcc aca tcc atc ttg ggc atc ggc act gtc ctt	4340
	Cys His Ser Thr Asp Ala Thr Ser Ile Leu Gly Ile Gly Thr Val Leu	
	1320 1325 1330	
25	gac caa gca gag act gcg ggg gcg aga ctg gtt gtg ctc gcc act gct	4388
	Asp Gln Ala Glu Thr Ala Gly Ala Arg Leu Val Val Leu Ala Thr Ala	
	1335 1340 1345	
30	acc cct ccg ggc tcc gtc act gtg tcc cat cct aac atc gag gag gtt	4436
	Thr Pro Pro Gly Ser Val Thr Val Ser His Pro Asn Ile Glu Glu Val	
	1350 1355 1360 1365	
35	gct ctg tcc acc acc gga gag atc ccc ttt tac ggc aag gct atc ccc	4484
	Ala Leu Ser Thr Thr Gly Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro	
	1370 1375 1380	
40	ctc gag gtg atc aag ggg gga aga cat ctc atc ttc tgc cac tca aag	4532
	Leu Glu Val Ile Lys Gly Gly Arg His Leu Ile Phe Cys His Ser Lys	
	1385 1390 1395	
45	aag aag tgc gac gag ctc gcc gcg aag ctg gtc gca ttg ggc atc aat	4580
	Lys Lys Cys Asp Glu Leu Ala Ala Lys Leu Val Ala Leu Gly Ile Asn	
	1400 1405 1410	
50	gcc gtg gcc tac tac cgc ggt ctt gac gtg tct gtc atc ccg acc agc	4628
	Ala Val Ala Tyr Tyr Arg Gly Leu Asp Val Ser Val Ile Pro Thr Ser	
	1415 1420 1425	
55	ggc gat gtt gtc gtc gtg tcg acc gat gct ctc atg act ggc ttt acc	4676
	Gly Asp Val Val Val Val Ser Thr Asp Ala Leu Met Thr Gly Phe Thr	
	1430 1435 1440 1445	
60	ggc gac ttc gac tct gtg ata gac tgc aac acg tgt gtc act cag aca	4724
	Gly Asp Phe Asp Ser Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr	
	1450 1455 1460	
65	gtc gat ttc agc ctt gac cct acc ttt acc att gag aca acc acg ctc	4772
	Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu	
	1465 1470 1475	
70	ccc cag gat gct gtc tcc agg act caa cgc cgg ggc agg act ggc agg	4820
	Pro Gln Asp Ala Val Ser Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg	
	1480 1485 1490	
75	ggg aag cca ggc atc tat aga ttt gtg gca ccg ggg gag cgc ccc tcc	4868
	Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro Gly Glu Arg Pro Ser	
	1495 1500 1505	
80	ggc atg ttc gac tcg tcc gtc ctc tgt gag tgc tat gac gcg ggc tgt	4916

	Gly Met Phe Asp Ser Ser Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys	
	1510 1515 1520 1525	
5	gct tgg tat gag ctc acg ccc gcc gag act aca gtt agg cta cga gcg Ala Trp Tyr Glu Leu Thr Pro Ala Glu Thr Val Arg Leu Arg Ala	4964
	1530 1535 1540	
10	tac atg aac acc ccg ggg ctt ccc gtg tgc cag gac cat ctt gaa ttt Tyr Met Asn Thr Pro Gly Leu Pro Val Cys Gln Asp His Leu Glu Phe	5012
	1545 1550 1555	
15	tgg gag ggc gtc ttt acg ggc ctc act cat ata gat gcc cac ttt tta Trp Glu Gly Val Phe Thr Gly Leu Thr His Ile Asp Ala His Phe Leu	5060
	1560 1565 1570	
20	tcc cag aca aag cag agt ggg gag aac ttt cct tac ctg gta gcg tac Ser Gln Thr Lys Gln Ser Gly Glu Asn Phe Pro Tyr Leu Val Ala Tyr	5108
	1575 1580 1585	
25	caa gcc acc gtg tgc gct agg gct caa gcc cct ccc cca tcg tgg gac Gln Ala Thr Val Cys Ala Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp	5156
	1590 1595 1600 1605	
30	cag atg tgg aag tgt ttg atc cgc ctt aaa ccc acc ctc cat ggg cca Gln Met Trp Lys Cys Leu Ile Arg Leu Lys Pro Thr Leu His Gly Pro	5204
	1610 1615 1620	
35	aca ccc ctg cta tac aga ctg ggc gct gtt cag aat gaa gtc acc ctg Thr Pro Leu Leu Tyr Arg Leu Gly Ala Val Gln Asn Glu Val Thr Leu	5252
	1625 1630 1635	
40	acg cac cca atc acc aaa tac atc atg aca tgc atg tcg gcc gac ctg Thr His Pro Ile Thr Lys Tyr Ile Met Thr Cys Met Ser Ala Asp Leu	5300
	1640 1645 1650	
45	gag gtc gtc acg agc acc tgg gtg ctc gtt ggc ggc gtc ctg gct gct Glu Val Val Thr Ser Thr Trp Val Leu Val Gly Gly Val Leu Ala Ala	5348
	1655 1660 1665	
50	ctg gcc gcg tat tgc ctg tca aca ggc tgc gtg gtc ata gtg ggc agg Leu Ala Ala Tyr Cys Leu Ser Thr Gly Cys Val Val Ile Val Gly Arg	5396
	1670 1675 1680 1685	
55	atc gtc ttg tcc ggg aag ccg gca att ata cct gac agg gag gtt ctc Ile Val Leu Ser Gly Lys Pro Ala Ile Ile Pro Asp Arg Glu Val Leu	5444
	1690 1695 1700	
60	tac cag gag ttc gat gag atg gaa gag tgc tct cag cac tta ccg tac Tyr Gln Glu Phe Asp Glu Met Glu Glu Cys Ser Gln His Leu Pro Tyr	5492
	1705 1710 1715	
65	atc gag caa ggg atg atg ctc gct gag cag ttc aag cag aag gcc ctc Ile Glu Gln Gly Met Met Leu Ala Glu Gln Phe Lys Gln Lys Ala Leu	5540
	1720 1725 1730	
70	ggc ctc ctg cag acc gcg tcc cgc cat gca gag gtt atc acc cct gct Gly Leu Leu Gln Thr Ala Ser Arg His Ala Glu Val Ile Thr Pro Ala	5588
	1735 1740 1745	
75	gtc cag acc aac tgg cag aaa ctc gag gtc ttt tgg gcg aag cac atg Val Gln Thr Asn Trp Gln Lys Leu Glu Val Phe Trp Ala Lys His Met	5636
	1750 1755 1760 1765	
80	tgg aat ttc atc agt ggg ata caa tac ttg gcg ggc ctg tca acg ctg Trp Asn Phe Ile Ser Gly Ile Gln Tyr Leu Ala Gly Leu Ser Thr Leu	5684
	1770 1775 1780	

	cct ggt aac ccc gcc att gct tca ttg atg gct ttt aca gct gcc gtc	5732
	Pro Gly Asn Pro Ala Ile Ala Ser Leu Met Ala Phe Thr Ala Ala Val	
	1785 1790 1795	
5	acc agc cca cta acc act ggc caa acc ctc ctc ttc aac ata ttg ggg	5780
	Thr Ser Pro Leu Thr Thr Gly Gln Thr Leu Leu Phe Asn Ile Leu Gly	
	1800 1805 1810	
10	ggg tgg gtg gct gcc cag ctc gcc gcc ccc ggt gcc gct act gcc ttt	5828
	Gly Trp Val Ala Ala Gln Leu Ala Ala Pro Gly Ala Ala Thr Ala Phe	
	1815 1820 1825	
15	gtg ggt gct ggc cta gct ggc gcc gcc atc ggc agc gtt gga ctg ggg	5876
	Val Gly Ala Gly Leu Ala Gly Ala Ala Ile Gly Ser Val Gly Leu Gly	
	1830 1835 1840 1845	
20	aag gtc ctc gtg gac att ctt gca ggg tat ggc gcg ggc gtg gcg gga	5924
	Lys Val Leu Val Asp Ile Leu Ala Gly Tyr Gly Ala Gly Val Ala Gly	
	1850 1855 1860	
25	gct ctt gta gca ttc aag atc atg agc ggt gag gtc ccc tcc acg gag	5972
	Ala Leu Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro Ser Thr Glu	
	1865 1870 1875	
30	gac ctg gtc aat ctg ctg ccc gcc atc ctc tcg cct gga gcc ctt gta	6020
	Asp Leu Val Asn Leu Leu Pro Ala Ile Leu Ser Pro Gly Ala Leu Val	
	1880 1885 1890	
35	gtc ggt gtg gtc tgc gca gca ata ctg cgc cgg cac gtt ggc ccg ggc	6068
	Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val Gly Pro Gly	
	1895 1900 1905	
40	gag ggg gca gtg caa tgg atg aac cgg cta ata gcc ttc gcc tcc cgg	6116
	Glu Gly Ala Val Gln Trp Met Asn Arg Leu Ile Ala Phe Ala Ser Arg	
	1910 1915 1920 1925	
45	ggg aac cat gtt tcc ccc acg cac tac gtg ccg gag agc gat gca gcc	6164
	Gly Asn His Val Ser Pro Thr His Tyr Val Pro Glu Ser Asp Ala Ala	
	1930 1935 1940	
50	gcc cgc gtc act gcc ata ctc agc agc ctc act gta acc cag ctc ctg	6212
	Ala Arg Val Thr Ala Ile Leu Ser Ser Leu Thr Val Thr Gln Leu Leu	
	1945 1950 1955	
55	agg cga ctg cat cag tgg ata agc tcg gag tgt acc act cca tgc tcc	6260
	Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys Thr Thr Pro Cys Ser	
	1960 1965 1970	
60	ggt tcc tgg cta agg gac atc tgg gac tgg ata tgc gag gtg ctg agc	6308
	Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp Ile Cys Glu Val Leu Ser	
	1975 1980 1985	
65	gac ttt aag acc tgg ctg aaa gcc aag ctc atg cca caa ctg cct ggg	6356
	Asp Phe Lys Thr Trp Leu Lys Ala Lys Leu Met Pro Gln Leu Pro Gly	
	1990 1995 2000 2005	
70	att ccc ttt gtg tcc tgc cag cgc ggg tat agg ggg gtc tgg cga gga	6404
	Ile Pro Phe Val Ser Cys Gln Arg Gly Tyr Arg Gly Val Trp Arg Gly	
	2010 2015 2020	
75	gac ggc att atg cac act cgc tgc cac tgt gga gct gag atc act gga	6452
	Asp Gly Ile Met His Thr Arg Cys His Cys Gly Ala Glu Ile Thr Gly	
	2025 2030 2035	
80	cat gtc aaa aac ggg acg atg agg atc gtc ggt cct agg acc tgc agg	6500

	His	Val	Lys	Asn	Gly	Thr	Met	Arg	Ile	Val	Gly	Pro	Arg	Thr	Cys	Arg	
		2040						2045					2050				
5	aac	atg	tgg	agt	ggg	acg	ttc	ccc	att	aac	gcc	tac	acc	acg	ggc	ccc	6548
	Asn	Met	Trp	Ser	Gly	Thr	Phe	Pro	Ile	Asn	Ala	Tyr	Thr	Thr	Gly	Pro	
		2055					2060					2065					
10	tgt	act	ccc	ctt	cct	gcg	ccg	aac	tat	aag	ttc	gcg	ctg	tgg	agg	gtg	6596
	Cys	Thr	Pro	Leu	Pro	Ala	Pro	Asn	Tyr	Lys	Phe	Ala	Leu	Trp	Arg	Val	
		2070				2075					2080				2085		
15	tct	gca	gag	gaa	tac	gtg	gag	ata	agg	cgg	gtg	ggg	gac	ttc	cac	tac	6644
	Ser	Ala	Glu	Glu	Tyr	Val	Glu	Ile	Arg	Arg	Val	Gly	Asp	Phe	His	Tyr	
					2090				2095					2100			
20	gta	tcg	ggt	atg	act	act	gac	aat	ctt	aaa	tgc	ccg	tgc	cag	atc	cca	6692
	Val	Ser	Gly	Met	Thr	Thr	Asp	Asn	Leu	Lys	Cys	Pro	Cys	Gln	Ile	Pro	
			2105					2110						2115			
25	tcg	ccc	gaa	ttt	ttc	aca	gaa	ttg	gac	ggg	gtg	cgc	cta	cac	agg	ttt	6740
	Ser	Pro	Glu	Phe	Phe	Thr	Glu	Leu	Asp	Gly	Val	Arg	Leu	His	Arg	Phe	
		2120					2125					2130					
30	gcg	ccc	cct	tgc	aag	ccc	ttg	ctg	cgg	gag	gag	gta	tca	ttc	aga	gta	6788
	Ala	Pro	Pro	Cys	Lys	Pro	Leu	Leu	Arg	Glu	Glu	Val	Ser	Phe	Arg	Val	
		2135					2140					2145					
35	gga	ctc	cac	gag	tac	ccg	gtg	ggg	tcg	caa	tta	cct	tgc	gag	ccc	gaa	6836
	Gly	Leu	His	Glu	Tyr	Pro	Val	Gly	Ser	Gln	Leu	Pro	Cys	Glu	Pro	Glu	
		2150				2155				2160					2165		
40	ccg	gac	gta	gcc	gtg	ttg	acg	tcc	atg	ctc	act	gat	ccc	tcc	cat	ata	6884
	Pro	Asp	Val	Ala	Val	Leu	Thr	Ser	Met	Leu	Thr	Asp	Pro	Ser	His	Ile	
					2170				2175					2180			
45	aca	gca	gag	gcg	gcc	ggg	aga	agg	ttg	gcg	aga	ggg	tca	ccc	cct	tct	6932
	Thr	Ala	Glu	Ala	Ala	Gly	Arg	Arg	Leu	Ala	Arg	Gly	Ser	Pro	Pro	Ser	
			2185				2190					2195					
50	atg	gcc	agc	tcc	tcg	gct	agc	cag	ctg	tcc	gct	cca	tct	ctc	aag	gca	6980
	Met	Ala	Ser	Ser	Ser	Ala	Ser	Gln	Leu	Ser	Ala	Pro	Ser	Leu	Lys	Ala	
		2200					2205					2210					
55	act	tgc	acc	gcc	aac	cat	gac	tcc	cct	gac	gcc	gag	ctc	ata	gag	gct	7028
	Thr	Cys	Thr	Ala	Asn	His	Asp	Ser	Pro	Asp	Ala	Glu	Leu	Ile	Glu	Ala	
		2215					2220					2225					
60	aac	ctc	ctg	tgg	agg	cag	gag	atg	ggc	ggc	aac	atc	acc	agg	gtt	gag	7076
	Asn	Leu	Leu	Trp	Arg	Gln	Glu	Met	Gly	Gly	Asn	Ile	Thr	Arg	Val	Glu	
		2230				2235					2240				2245		
65	tca	gag	aac	aaa	gtg	gtg	att	ctg	gac	tcc	ttc	gat	ccg	ctt	gtg	gca	7124
	Ser	Glu	Asn	Lys	Val	Val	Ile	Leu	Asp	Ser	Phe	Asp	Pro	Leu	Val	Ala	
					2250				2255					2260			
70	gag	gag	gat	gag	cgg	gag	gtc	tcc	gta	cct	gca	gaa	att	ctg	cgg	aag	7172
	Glu	Glu	Asp	Glu	Arg	Glu	Val	Ser	Val	Pro	Ala	Glu	Ile	Leu	Arg	Lys	
					2265				2270					2275			
75	tct	cgg	aga	ttc	gcc	cgg	gcc	ctg	ccc	gtc	tgg	gcg	cgg	ccg	gac	tac	7220
	Ser	Arg	Arg	Phe	Ala	Arg	Ala	Leu	Pro	Val	Trp	Ala	Arg	Pro	Asp	Tyr	
		2280					2285					2290					
80	aac	ccc	ccg	cta	gta	gag	acg	tgg	aaa	aag	cct	gac	tac	gaa	cca	cct	7268
	Asn	Pro	Pro	Leu	Val	Glu	Thr	Trp	Lys	Lys	Pro	Asp	Tyr	Glu	Pro	Pro	
		2295					2300					2305					

5	gtg gtc cat ggc tgc ccg cta cca cct cca cgg tcc cct cct gtg cct Val Val His Gly Cys Pro Leu Pro Pro Pro Arg Ser Pro Pro Val Pro 2310 2315 2320 2325	7316
	ccg cct cgg aaa aag cgt acg gtg gtc ctc acc gaa tca acc cta tct Pro Pro Arg Lys Lys Arg Thr Val Val Leu Thr Glu Ser Thr Leu Ser 2330 2335 2340	7364
10	act gcc ttg gcc gag ctt gcc acc aaa agt ttt ggc agc tcc tca act Thr Ala Leu Ala Glu Leu Ala Thr Lys Ser Phe Gly Ser Ser Ser Thr 2345 2350 2355	7412
15	tcc ggc att acg ggc gac aat acg aca aca tcc tct gag ccc gcc cct Ser Gly Ile Thr Gly Asp Asn Thr Thr Ser Ser Glu Pro Ala Pro 2360 2365 2370	7460
20	tct ggc tgc ccc ccc gac tcc gac gtt gag tcc tat tct tcc atg ccc Ser Gly Cys Pro Pro Asp Ser Asp Val Glu Ser Tyr Ser Ser Met Pro 2375 2380 2385	7508
25	ccc ctg gag ggg gag cct ggg gat ccg gat ctc agc gac ggg tca tgg Pro Leu Glu Gly Glu Pro Gly Asp Pro Asp Leu Ser Asp Gly Ser Trp 2390 2395 2400 2405	7556
	tcg acg gtc agt agt ggg gcc gac acg gaa gat gtc gtg tgc tgc tca Ser Thr Val Ser Ser Gly Ala Asp Thr Glu Asp Val Val Cys Cys Ser 2410 2415 2420	7604
30	atg tct tat tcc tgg aca ggc gca ctc gtc acc ccg tgc gct gcg gaa Met Ser Tyr Ser Trp Thr Gly Ala Leu Val Thr Pro Cys Ala Ala Glu 2425 2430 2435	7652
35	gaa caa aaa ctg ccc atc aac gca ctg agc aac tcg ttg cta cgc cat Glu Gln Lys Leu Pro Ile Asn Ala Leu Ser Asn Ser Leu Leu Arg His 2440 2445 2450	7700
40	cac aat ctg gtg tat tcc acc act tca cgc agt gct tgc caa agg cag His Asn Leu Val Tyr Ser Thr Thr Ser Arg Ser Ala Cys Gln Arg Gln 2455 2460 2465	7748
45	aag aaa gtc aca ttt gac aga ctg caa gtt ctg gac agc cat tac cag Lys Lys Val Thr Phe Asp Arg Leu Gln Val Leu Asp Ser His Tyr Gln 2470 2475 2480 2485	7796
	gac gtg ctc aag gag gtc aaa gca gcg gcg tca aaa gtg aag gct aac Asp Val Leu Lys Glu Val Lys Ala Ala Ala Ser Lys Val Lys Ala Asn 2490 2495 2500	7844
50	ttg cta tcc gta gag gaa gct tgc agc ctg acg ccc cca cat tca gcc Leu Leu Ser Val Glu Glu Ala Cys Ser Leu Thr Pro Pro His Ser Ala 2505 2510 2515	7892
55	aaa tcc aag ttt ggc tat ggg gca aaa gac gtc cgt tgc cat gcc aga Lys Ser Lys Phe Gly Tyr Gly Ala Lys Asp Val Arg Cys His Ala Arg 2520 2525 2530	7940
60	aag gcc gta gcc cac atc aac tcc gtg tgg aaa gac ctt ctg gaa gac Lys Ala Val Ala His Ile Asn Ser Val Trp Lys Asp Leu Leu Glu Asp 2535 2540 2545	7988
65	agt gta aca cca ata gac act acc atc atg gcc aag aac gag gtt ttc Ser Val Thr Pro Ile Asp Thr Thr Ile Met Ala Lys Asn Glu Val Phe 2550 2555 2560 2565	8036
	tgc gtt cag cct gag aag ggg ggt cgt aag cca gct cgt ctc atc gtg	8084

	Cys	Val	Gln	Pro	Glu	Lys	Gly	Gly	Arg	Lys	Pro	Ala	Arg	Leu	Ile	Val	
					2570					2575					2580		
5	ttc	ccc	gac	ctg	ggc	gtg	cgc	gtg	tgc	gag	aag	atg	gcc	ctg	tac	gac	8132
	Phe	Pro	Asp	Leu	Gly	Val	Arg	Val	Cys	Glu	Lys	Met	Ala	Leu	Tyr	Asp	
				2585					2590					2595			
10	gtg	gtt	agc	aag	ctc	ccc	ctg	gcc	gtg	atg	gga	agc	tcc	tac	gga	ttc	8180
	Val	Val	Ser	Lys	Leu	Pro	Leu	Ala	Val	Met	Gly	Ser	Ser	Tyr	Gly	Phe	
			2600					2605					2610				
15	caa	tac	tca	cca	gga	cag	cgg	gtt	gaa	ttc	ctc	gtg	caa	gcg	tgg	aag	8228
	Gln	Tyr	Ser	Pro	Gly	Gln	Arg	Val	Glu	Phe	Leu	Val	Gln	Ala	Trp	Lys	
		2615					2620					2625					
	tcc	aag	aag	acc	ccg	atg	ggg	ttc	tgc	tat	gat	acc	cgc	tgt	ttt	gac	8276
	Ser	Lys	Lys	Thr	Pro	Met	Gly	Phe	Ser	Tyr	Asp	Thr	Arg	Cys	Phe	Asp	
		2630				2635					2640				2645		
20	tcc	aca	gtc	act	gag	agc	gac	atc	cgt	acg	gag	gag	gca	att	tac	caa	8324
	Ser	Thr	Val	Thr	Glu	Ser	Asp	Ile	Arg	Thr	Glu	Glu	Ala	Ile	Tyr	Gln	
				2650						2655					2660		
25	tgt	tgt	gac	ctg	gac	ccc	caa	gcc	cgc	gtg	gcc	atc	aag	tcc	ctc	act	8372
	Cys	Cys	Asp	Leu	Asp	Pro	Gln	Ala	Arg	Val	Ala	Ile	Lys	Ser	Leu	Thr	
			2665					2670					2675				
30	gag	agg	ctt	tat	gtt	ggg	ggc	cct	ctt	acc	aat	tca	agg	ggg	gaa	aac	8420
	Glu	Arg	Leu	Tyr	Val	Gly	Gly	Pro	Leu	Thr	Asn	Ser	Arg	Gly	Glu	Asn	
		2680					2685					2690					
35	tgc	ggc	tac	cgc	agg	tgc	cgc	gcg	agc	ggc	gta	ctg	aca	act	agc	tgt	8468
	Cys	Gly	Tyr	Arg	Arg	Cys	Arg	Ala	Ser	Gly	Val	Leu	Thr	Thr	Ser	Cys	
		2695					2700				2705						
	ggt	aac	acc	ctc	act	tgc	tac	atc	aag	gcc	cgg	gca	gcc	tgt	cga	gcc	8516
	Gly	Asn	Thr	Leu	Thr	Cys	Tyr	Ile	Lys	Ala	Arg	Ala	Ala	Cys	Arg	Ala	
		2710				2715				2720					2725		
40	gca	ggg	ctc	cag	gac	tgc	acc	atg	ctc	gtg	tgt	ggc	gac	gac	tta	gtc	8564
	Ala	Gly	Leu	Gln	Asp	Cys	Thr	Met	Leu	Val	Cys	Gly	Asp	Asp	Leu	Val	
				2730						2735					2740		
45	gtt	atc	tgt	gaa	agt	gcg	ggg	gtc	cag	gag	gac	gcg	gcg	agc	ctg	aga	8612
	Val	Ile	Cys	Glu	Ser	Ala	Gly	Val	Gln	Glu	Asp	Ala	Ala	Ser	Leu	Arg	
			2745					2750						2755			
50	gcc	ttc	acg	gag	gct	atg	acc	agg	tac	tcc	gcc	ccc	ccc	ggg	gac	ccc	8660
	Ala	Phe	Thr	Glu	Ala	Met	Thr	Arg	Tyr	Ser	Ala	Pro	Pro	Gly	Asp	Pro	
		2760						2765					2770				
55	cca	caa	cca	gaa	tac	gac	ttg	gag	ctt	ata	aca	tca	tgc	tcc	tcc	aac	8708
	Pro	Gln	Pro	Glu	Tyr	Asp	Leu	Glu	Leu	Ile	Thr	Ser	Cys	Ser	Ser	Asn	
		2775					2780					2785					
	gtg	tca	gtc	gcc	cac	gac	ggc	gct	gga	aag	agg	gtc	tac	tac	ctt	acc	8756
	Val	Ser	Val	Ala	His	Asp	Gly	Ala	Gly	Lys	Arg	Val	Tyr	Tyr	Leu	Thr	
		2790				2795				2800					2805		
60	cgt	gac	cct	aca	acc	ccc	ctc	gcg	aga	gcc	gcg	tgg	gag	aca	gca	aga	8804
	Arg	Asp	Pro	Thr	Thr	Pro	Leu	Ala	Arg	Ala	Ala	Trp	Glu	Thr	Ala	Arg	
				2810					2815						2820		
65	cac	act	cca	gtc	aat	tcc	tgg	cta	ggc	aac	ata	atc	atg	ttt	gcc	ccc	8852
	His	Thr	Pro	Val	Asn	Ser	Trp	Leu	Gly	Asn	Ile	Ile	Met	Phe	Ala	Pro	
			2825					2830						2835			

	aca ctg tgg gcg agg atg ata ctg atg acc cat ttc ttt agc gtc ctc	8900
	Thr Leu Trp Ala Arg Met Ile Leu Met Thr His Phe Phe Ser Val Leu	
	2840 2845 2850	
5	ata gcc agg gat cag ctt gaa cag gct ctt aac tgt gag atc tac gga	8948
	Ile Ala Arg Asp Gln Leu Glu Gln Ala Leu Asn Cys Glu Ile Tyr Gly	
	2855 2860 2865	
10	gcc tgc tac tcc ata gaa cca ctg gat cta cct cca atc att caa aga	8996
	Ala Cys Tyr Ser Ile Glu Pro Leu Asp Leu Pro Pro Ile Ile Gln Arg	
	2870 2875 2880 2885	
15	ctc cat ggc ctc agc gca ttt tca ctc cac agt tac tct cca ggt gaa	9044
	Leu His Gly Leu Ser Ala Phe Ser Leu His Ser Tyr Ser Pro Gly Glu	
	2890 2895 2900	
20	atc aat agg gtg gcc gca tgc ctc aga aaa ctt ggg gtc ccg ccc ttg	9092
	Ile Asn Arg Val Ala Ala Cys Leu Arg Lys Leu Gly Val Pro Pro Leu	
	2905 2910 2915	
25	cga gct tgg aga cac cgg gcc cgg agc gtc cgc gct agg ctt ctg tcc	9140
	Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg Ala Arg Leu Leu Ser	
	2920 2925 2930	
30	aga gga ggc agg gct gcc ata tgt ggc aag tac ctc ttc aac tgg gca	9188
	Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr Leu Phe Asn Trp Ala	
	2935 2940 2945	
35	gta aga aca aag ctc aaa ctc act cca ata gcg gcc gct ggc cgg ctg	9236
	Val Arg Thr Lys Leu Lys Leu Thr Pro Ile Ala Ala Ala Gly Arg Leu	
	2950 2955 2960 2965	
40	gac ttg tcc ggt tgg ttc acg gct ggc tac agc ggg gga gac att tat	9284
	Asp Leu Ser Gly Trp Phe Thr Ala Gly Tyr Ser Gly Gly Asp Ile Tyr	
	2970 2975 2980	
45	cac agc gtg tct cat gcc cgg ccc cgc tgg ttc tgg ttt tgc cta ctc	9332
	His Ser Val Ser His Ala Arg Pro Arg Trp Phe Trp Phe Cys Leu Leu	
	2985 2990 2995	
50	ctg ctc gct gca ggg gta ggc atc tac ctc ctc ccc aac cga tga	9377
	Leu Leu Ala Ala Gly Val Gly Ile Tyr Leu Leu Pro Asn Arg	
	3000 3005 3010	
55	agggtggggg aaacactccg gcctcttaag ccatttctctg tttttttttt tttttttttt	9437
	ttttttttct tttttttttt ctttcctttc cttctttttt tcttttcttt ttcccttctt	9497
60	taatggtggc tccatcttag ccctagtcac ggctagctgt gaaagggtccg tgagccgcat	9557
	gactgcagag agtgcgtgata ctggcctctc tgcagatcat gt	9599
65	<210> 14	
	<211> 3011	
	<212> PRT	
	<213> Hepatitis C virus	
60	<400> 14	
	Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn	
	1 5 10 15	
	Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly	
	20 25 30	
65	Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala	
	35 40 45	

	Thr	Arg	Lys	Thr	Ser	Glu	Arg	Ser	Gln	Pro	Arg	Gly	Arg	Arg	Gln	Pro
	50						55					60				
	Ile	Pro	Lys	Ala	Arg	Arg	Pro	Glu	Gly	Arg	Thr	Trp	Ala	Gln	Pro	Gly
	65					70					75				80	
5	Tyr	Pro	Trp	Pro	Leu	Tyr	Gly	Asn	Glu	Gly	Cys	Gly	Trp	Ala	Gly	Trp
					85					90					95	
	Leu	Leu	Ser	Pro	Arg	Gly	Ser	Arg	Pro	Ser	Trp	Gly	Pro	Thr	Asp	Pro
				100					105					110		
10	Arg	Arg	Arg	Ser	Arg	Asn	Leu	Gly	Lys	Val	Ile	Asp	Thr	Leu	Thr	Cys
				115				120					125			
	Gly	Phe	Ala	Asp	Leu	Met	Gly	Tyr	Ile	Pro	Leu	Val	Gly	Ala	Pro	Leu
		130					135					140				
	Gly	Gly	Ala	Ala	Arg	Ala	Leu	Ala	His	Gly	Val	Arg	Val	Leu	Glu	Asp
		145				150					155				160	
15	Gly	Val	Asn	Tyr	Ala	Thr	Gly	Asn	Leu	Pro	Gly	Cys	Ser	Phe	Ser	Ile
					165					170					175	
	Phe	Leu	Leu	Ala	Leu	Leu	Ser	Cys	Leu	Thr	Val	Pro	Ala	Ser	Ala	Tyr
				180					185						190	
20	Gln	Val	Arg	Asn	Ser	Ser	Gly	Leu	Tyr	His	Val	Thr	Asn	Asp	Cys	Pro
			195					200					205			
	Asn	Ser	Ser	Ile	Val	Tyr	Glu	Ala	Ala	Asp	Ala	Ile	Leu	His	Thr	Pro
		210					215					220				
	Gly	Cys	Val	Pro	Cys	Val	Arg	Glu	Gly	Asn	Ala	Ser	Arg	Cys	Trp	Val
		225				230					235				240	
25	Ala	Val	Thr	Pro	Thr	Val	Ala	Thr	Arg	Asp	Gly	Lys	Leu	Pro	Thr	Thr
					245					250					255	
	Gln	Leu	Arg	Arg	His	Ile	Asp	Leu	Leu	Val	Gly	Ser	Ala	Thr	Leu	Cys
				260					265					270		
30	Ser	Ala	Leu	Tyr	Val	Gly	Asp	Leu	Cys	Gly	Ser	Val	Phe	Leu	Val	Gly
			275					280					285			
	Gln	Leu	Phe	Thr	Phe	Ser	Pro	Arg	Arg	His	Trp	Thr	Thr	Gln	Asp	Cys
		290					295					300				
	Asn	Cys	Ser	Ile	Tyr	Pro	Gly	His	Ile	Thr	Gly	His	Arg	Met	Ala	Trp
		305				310					315					320
35	Asp	Met	Met	Met	Asn	Trp	Ser	Pro	Thr	Ala	Ala	Leu	Val	Val	Ala	Gln
					325					330					335	
	Leu	Leu	Arg	Ile	Pro	Gln	Ala	Ile	Met	Asp	Met	Ile	Ala	Gly	Ala	His
				340					345					350		
40	Trp	Gly	Val	Leu	Ala	Gly	Ile	Ala	Tyr	Phe	Ser	Met	Val	Gly	Asn	Trp
			355					360					365			
	Ala	Lys	Val	Leu	Val	Val	Leu	Leu	Phe	Ala	Gly	Val	Asp	Ala	Glu	
		370					375				380					
	Thr	His	Val	Thr	Gly	Gly	Asn	Ala	Gly	Arg	Thr	Thr	Ala	Gly	Leu	Val
		385				390					395				400	
45	Gly	Leu	Leu	Thr	Pro	Gly	Ala	Lys	Gln	Asn	Ile	Gln	Leu	Ile	Asn	Thr
					405					410					415	
	Asn	Gly	Ser	Trp	His	Ile	Asn	Ser	Thr	Ala	Leu	Asn	Cys	Asn	Glu	Ser
				420					425					430		
50	Leu	Asn	Thr	Gly	Trp	Leu	Ala	Gly	Leu	Phe	Tyr	Gln	His	Lys	Phe	Asn
			435					440					445			
	Ser	Ser	Gly	Cys	Pro	Glu	Arg	Leu	Ala	Ser	Cys	Arg	Arg	Leu	Thr	Asp
			450				455					460				
	Phe	Ala	Gln	Gly	Trp	Gly	Pro	Ile	Ser	Tyr	Ala	Asn	Gly	Ser	Gly	Leu
		465				470					475				480	
55	Asp	Glu	Arg	Pro	Tyr	Cys	Trp	His	Tyr	Pro	Pro	Arg	Pro	Cys	Gly	Ile
					485					490					495	
	Val	Pro	Ala	Lys	Ser	Val	Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	Ser
				500					505					510		
60	Pro	Val	Val	Val	Gly	Thr	Thr	Asp	Arg	Ser	Gly	Ala	Pro	Thr	Tyr	Ser
			515					520					525			
	Trp	Gly	Ala	Asn	Asp	Thr	Asp	Val	Phe	Val	Leu	Asn	Asn	Thr	Arg	Pro
		530					535					540				
	Pro	Leu	Gly	Asn	Trp	Phe	Gly	Cys	Thr	Trp	Met	Asn	Ser	Thr	Gly	Phe
		545				550					555				560	
65	Thr	Lys	Val	Cys	Gly	Ala	Pro	Pro	Cys	Val	Ile	Gly	Gly	Val	Gly	Asn
					565					570					575	

	Asn	Thr	Leu	Leu	Cys	Pro	Thr	Asp	Cys	Phe	Arg	Lys	His	Pro	Glu	Ala
			580						585					590		
	Thr	Tyr	Ser	Arg	Cys	Gly	Ser	Gly	Pro	Trp	Ile	Thr	Pro	Arg	Cys	Met
			595					600					605			
5	Val	Asp	Tyr	Pro	Tyr	Arg	Leu	Trp	His	Tyr	Pro	Cys	Thr	Ile	Asn	Tyr
		610					615					620				
	Thr	Ile	Phe	Lys	Val	Arg	Met	Tyr	Val	Gly	Gly	Val	Glu	His	Arg	Leu
		625				630					635					640
10	Glu	Ala	Ala	Cys	Asn	Trp	Thr	Arg	Gly	Glu	Arg	Cys	Asp	Leu	Glu	Asp
					645					650					655	
	Arg	Asp	Arg	Ser	Glu	Leu	Ser	Pro	Leu	Leu	Leu	Ser	Thr	Thr	Gln	Trp
				660					665					670		
	Gln	Val	Leu	Pro	Cys	Ser	Phe	Thr	Thr	Leu	Pro	Ala	Leu	Ser	Thr	Gly
			675					680					685			
15	Leu	Ile	His	Leu	His	Gln	Asn	Ile	Val	Asp	Val	Gln	Tyr	Leu	Tyr	Gly
		690					695					700				
	Val	Gly	Ser	Ser	Ile	Ala	Ser	Trp	Ala	Ile	Lys	Trp	Glu	Tyr	Val	Val
		705				710					715					720
20	Leu	Leu	Phe	Leu	Leu	Leu	Ala	Asp	Ala	Arg	Val	Cys	Ser	Cys	Leu	Trp
					725				730						735	
	Met	Met	Leu	Leu	Ile	Ser	Gln	Ala	Glu	Ala	Ala	Leu	Glu	Asn	Leu	Val
				740					745					750		
	Ile	Leu	Asn	Ala	Ala	Ser	Leu	Ala	Gly	Thr	His	Gly	Leu	Val	Ser	Phe
			755					760					765			
25	Leu	Val	Phe	Phe	Cys	Phe	Ala	Trp	Tyr	Leu	Lys	Gly	Arg	Trp	Val	Pro
		770					775					780				
	Gly	Ala	Val	Tyr	Ala	Leu	Tyr	Gly	Met	Trp	Pro	Leu	Leu	Leu	Leu	Leu
		785				790					795					800
30	Leu	Ala	Leu	Pro	Gln	Arg	Ala	Tyr	Ala	Leu	Asp	Thr	Glu	Val	Ala	Ala
					805					810					815	
	Ser	Cys	Gly	Gly	Val	Val	Leu	Val	Gly	Leu	Met	Ala	Leu	Thr	Leu	Ser
				820					825					830		
	Pro	Tyr	Tyr	Lys	Arg	Tyr	Ile	Ser	Trp	Cys	Met	Trp	Trp	Leu	Gln	Tyr
			835					840						845		
35	Phe	Leu	Thr	Arg	Val	Glu	Ala	Gln	Leu	His	Val	Trp	Val	Pro	Pro	Leu
		850					855					860				
	Asn	Val	Arg	Gly	Gly	Arg	Asp	Ala	Val	Ile	Leu	Leu	Met	Cys	Val	Val
		865				870					875					880
40	His	Pro	Thr	Leu	Val	Phe	Asp	Ile	Thr	Lys	Leu	Leu	Leu	Ala	Ile	Phe
					885					890					895	
	Gly	Pro	Leu	Trp	Ile	Leu	Gln	Ala	Ser	Leu	Leu	Lys	Val	Pro	Tyr	Phe
				900					905					910		
	Val	Arg	Val	Gln	Gly	Leu	Leu	Arg	Ile	Cys	Ala	Leu	Ala	Arg	Lys	Ile
			915					920					925			
45	Ala	Gly	Gly	His	Tyr	Val	Gln	Met	Ala	Ile	Ile	Lys	Leu	Gly	Ala	Leu
		930					935					940				
	Thr	Gly	Thr	Tyr	Val	Tyr	Asn	His	Leu	Thr	Pro	Leu	Arg	Asp	Trp	Ala
		945				950					955					960
50	His	Asn	Gly	Leu	Arg	Asp	Leu	Ala	Val	Ala	Val	Glu	Pro	Val	Val	Phe
					965					970					975	
	Ser	Arg	Met	Glu	Thr	Lys	Leu	Ile	Thr	Trp	Gly	Ala	Asp	Thr	Ala	Ala
				980					985					990		
	Cys	Gly	Asp	Ile	Ile	Asn	Gly	Leu	Pro	Val	Ser	Ala	Arg	Arg	Gly	Gln
			995				1000					1005				
55	Glu	Ile	Leu	Leu	Gly	Pro	Ala	Asp	Gly	Met	Val	Ser	Lys	Gly	Trp	Arg
		1010					1015					1020				
	Leu	Leu	Ala	Pro	Ile	Thr	Ala	Tyr	Ala	Gln	Gln	Thr	Arg	Gly	Leu	Leu
		1025				1030				1035						1040
60	Gly	Cys	Ile	Ile	Thr	Ser	Leu	Thr	Gly	Arg	Asp	Lys	Asn	Gln	Val	Glu
					1045				1050					1055		
	Gly	Glu	Val	Gln	Ile	Val	Ser	Thr	Ala	Thr	Gln	Thr	Phe	Leu	Ala	Thr
			1060					1065					1070			
	Cys	Ile	Asn	Gly	Val	Cys	Trp	Thr	Val	Tyr	His	Gly	Ala	Gly	Thr	Arg
		1075					1080					1085				
65	Thr	Ile	Ala	Ser	Pro	Lys	Gly	Pro	Val	Ile	Gln	Met	Tyr	Thr	Asn	Val
		1090					1095					1100				

Asp Gln Asp Leu Val Gly Trp Pro Ala Pro Gln Gly Ser Arg Ser Leu
 1105 1110 1115 1120
 Thr Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr Leu Val Thr Arg His
 1125 1130 1135
 5 Ala Asp Val Ile Pro Val Arg Arg Arg Gly Asp Ser Arg Gly Ser Leu
 1140 1145 1150
 Leu Ser Pro Arg Pro Ile Ser Tyr Leu Lys Gly Ser Ser Gly Gly Pro
 1155 1160 1165
 10 Leu Leu Cys Pro Ala Gly His Ala Val Gly Leu Phe Arg Ala Ala Val
 1170 1175 1180
 Cys Thr Arg Gly Val Ala Lys Ala Val Asp Phe Ile Pro Val Glu Asn
 1185 1190 1195 1200
 Leu Gly Thr Thr Met Arg Ser Pro Val Phe Thr Asp Asn Ser Ser Pro
 1205 1210 1215
 15 Pro Ala Val Pro Gln Ser Phe Gln Val Ala His Leu His Ala Pro Thr
 1220 1225 1230
 Gly Ser Gly Lys Ser Thr Lys Val Pro Ala Ala Tyr Ala Ala Gln Gly
 1235 1240 1245
 20 Tyr Lys Val Leu Val Leu Asn Pro Ser Val Ala Ala Thr Leu Gly Phe
 1250 1255 1260
 Gly Ala Tyr Met Ser Lys Ala His Gly Val Asp Pro Asn Ile Arg Thr
 1265 1270 1275 1280
 Gly Val Arg Thr Ile Thr Thr Gly Ser Pro Ile Thr Tyr Ser Thr Tyr
 1285 1290 1295
 25 Gly Lys Phe Leu Ala Asp Gly Gly Cys Ser Gly Gly Ala Tyr Asp Ile
 1300 1305 1310
 Ile Ile Cys Asp Glu Cys His Ser Thr Asp Ala Thr Ser Ile Leu Gly
 1315 1320 1325
 30 Ile Gly Thr Val Leu Asp Gln Ala Glu Thr Ala Gly Ala Arg Leu Val
 1330 1335 1340
 Val Leu Ala Thr Ala Thr Pro Pro Gly Ser Val Thr Val Ser His Pro
 1345 1350 1355 1360
 Asn Ile Glu Glu Val Ala Leu Ser Thr Thr Gly Glu Ile Pro Phe Tyr
 1365 1370 1375
 35 Gly Lys Ala Ile Pro Leu Glu Val Ile Lys Gly Gly Arg His Leu Ile
 1380 1385 1390
 Phe Cys His Ser Lys Lys Lys Cys Asp Glu Leu Ala Ala Lys Leu Val
 1395 1400 1405
 40 Ala Leu Gly Ile Asn Ala Val Ala Tyr Tyr Arg Gly Leu Asp Val Ser
 1410 1415 1420
 Val Ile Pro Thr Ser Gly Asp Val Val Val Val Ser Thr Asp Ala Leu
 1425 1430 1435 1440
 Met Thr Gly Phe Thr Gly Asp Phe Asp Ser Val Ile Asp Cys Asn Thr
 1445 1450 1455
 45 Cys Val Thr Gln Thr Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile
 1460 1465 1470
 Glu Thr Thr Thr Leu Pro Gln Asp Ala Val Ser Arg Thr Gln Arg Arg
 1475 1480 1485
 50 Gly Arg Thr Gly Arg Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro
 1490 1495 1500
 Gly Glu Arg Pro Ser Gly Met Phe Asp Ser Ser Val Leu Cys Glu Cys
 1505 1510 1515 1520
 Tyr Asp Ala Gly Cys Ala Trp Tyr Glu Leu Thr Pro Ala Glu Thr Thr
 1525 1530 1535
 55 Val Arg Leu Arg Ala Tyr Met Asn Thr Pro Gly Leu Pro Val Cys Gln
 1540 1545 1550
 Asp His Leu Glu Phe Trp Glu Gly Val Phe Thr Gly Leu Thr His Ile
 1555 1560 1565
 60 Asp Ala His Phe Leu Ser Gln Thr Lys Gln Ser Gly Glu Asn Phe Pro
 1570 1575 1580
 Tyr Leu Val Ala Tyr Gln Ala Thr Val Cys Ala Arg Ala Gln Ala Pro
 1585 1590 1595 1600
 Pro Pro Ser Trp Asp Gln Met Trp Lys Cys Leu Ile Arg Leu Lys Pro
 1605 1610 1615
 65 Thr Leu His Gly Pro Thr Pro Leu Leu Tyr Arg Leu Gly Ala Val Gln
 1620 1625 1630

	Asn	Glu	Val	Thr	Leu	Thr	His	Pro	Ile	Thr	Lys	Tyr	Ile	Met	Thr	Cys	
		1635						1640					1645				
	Met	Ser	Ala	Asp	Leu	Glu	Val	Val	Thr	Ser	Thr	Trp	Val	Leu	Val	Gly	
		1650					1655					1660					
5	Gly	Val	Leu	Ala	Ala	Leu	Ala	Ala	Tyr	Cys	Leu	Ser	Thr	Gly	Cys	Val	
		1665				1670					1675					1680	
	Val	Ile	Val	Gly	Arg	Ile	Val	Leu	Ser	Gly	Lys	Pro	Ala	Ile	Ile	Pro	
				1685						1690					1695		
10	Asp	Arg	Glu	Val	Leu	Tyr	Gln	Glu	Phe	Asp	Glu	Met	Glu	Glu	Cys	Ser	
			1700						1705					1710			
	Gln	His	Leu	Pro	Tyr	Ile	Glu	Gln	Gly	Met	Met	Leu	Ala	Glu	Gln	Phe	
		1715					1720						1725				
	Lys	Gln	Lys	Ala	Leu	Gly	Leu	Leu	Gln	Thr	Ala	Ser	Arg	His	Ala	Glu	
		1730				1735					1740						
15	Val	Ile	Thr	Pro	Ala	Val	Gln	Thr	Asn	Trp	Gln	Lys	Leu	Glu	Val	Phe	
		1745			1750					1755					1760		
	Trp	Ala	Lys	His	Met	Trp	Asn	Phe	Ile	Ser	Gly	Ile	Gln	Tyr	Leu	Ala	
				1765					1770					1775			
20	Gly	Leu	Ser	Thr	Leu	Pro	Gly	Asn	Pro	Ala	Ile	Ala	Ser	Leu	Met	Ala	
		1780						1785					1790				
	Phe	Thr	Ala	Ala	Val	Thr	Ser	Pro	Leu	Thr	Thr	Gly	Gln	Thr	Leu	Leu	
		1795					1800						1805				
	Phe	Asn	Ile	Leu	Gly	Gly	Trp	Val	Ala	Ala	Gln	Leu	Ala	Ala	Pro	Gly	
		1810				1815					1820						
25	Ala	Ala	Thr	Ala	Phe	Val	Gly	Ala	Gly	Leu	Ala	Gly	Ala	Ala	Ile	Gly	
		1825				1830					1835				1840		
	Ser	Val	Gly	Leu	Gly	Lys	Val	Leu	Val	Asp	Ile	Leu	Ala	Gly	Tyr	Gly	
				1845					1850					1855			
30	Ala	Gly	Val	Ala	Gly	Ala	Leu	Val	Ala	Phe	Lys	Ile	Met	Ser	Gly	Glu	
		1860						1865					1870				
	Val	Pro	Ser	Thr	Glu	Asp	Leu	Val	Asn	Leu	Leu	Pro	Ala	Ile	Leu	Ser	
		1875					1880					1885					
	Pro	Gly	Ala	Leu	Val	Val	Gly	Val	Val	Cys	Ala	Ala	Ile	Leu	Arg	Arg	
		1890				1895						1900					
35	His	Val	Gly	Pro	Gly	Glu	Gly	Ala	Val	Gln	Trp	Met	Asn	Arg	Leu	Ile	
		1905				1910				1915					1920		
	Ala	Phe	Ala	Ser	Arg	Gly	Asn	His	Val	Ser	Pro	Thr	His	Tyr	Val	Pro	
				1925				1930					1935				
40	Glu	Ser	Asp	Ala	Ala	Arg	Val	Thr	Ala	Ile	Leu	Ser	Ser	Leu	Thr		
		1940					1945					1950					
	Val	Thr	Gln	Leu	Leu	Arg	Arg	Leu	His	Gln	Trp	Ile	Ser	Ser	Glu	Cys	
		1955					1960					1965					
	Thr	Thr	Pro	Cys	Ser	Gly	Ser	Trp	Leu	Arg	Asp	Ile	Trp	Asp	Trp	Ile	
		1970				1975					1980						
45	Cys	Glu	Val	Leu	Ser	Asp	Phe	Lys	Thr	Trp	Leu	Lys	Ala	Lys	Leu	Met	
		1985				1990				1995					2000		
	Pro	Gln	Leu	Pro	Gly	Ile	Pro	Phe	Val	Ser	Cys	Gln	Arg	Gly	Tyr	Arg	
				2005				2010					2015				
50	Gly	Val	Trp	Arg	Gly	Asp	Gly	Ile	Met	His	Thr	Arg	Cys	His	Cys	Gly	
		2020						2025					2030				
	Ala	Glu	Ile	Thr	Gly	His	Val	Lys	Asn	Gly	Thr	Met	Arg	Ile	Val	Gly	
		2035					2040					2045					
	Pro	Arg	Thr	Cys	Arg	Asn	Met	Trp	Ser	Gly	Thr	Phe	Pro	Ile	Asn	Ala	
		2050				2055					2060						
55	Tyr	Thr	Thr	Gly	Pro	Cys	Thr	Pro	Leu	Pro	Ala	Pro	Asn	Tyr	Lys	Phe	
		2065				2070				2075					2080		
	Ala	Leu	Trp	Arg	Val	Ser	Ala	Glu	Glu	Tyr	Val	Glu	Ile	Arg	Arg	Val	
				2085				2090					2095				
60	Gly	Asp	Phe	His	Tyr	Val	Ser	Gly	Met	Thr	Thr	Asp	Asn	Leu	Lys	Cys	
		2100						2105				2110					
	Pro	Cys	Gln	Ile	Pro	Ser	Pro	Glu	Phe	Phe	Thr	Glu	Leu	Asp	Gly	Val	
		2115					2120					2125					
	Arg	Leu	His	Arg	Phe	Ala	Pro	Pro	Cys	Lys	Pro	Leu	Leu	Arg	Glu	Glu	
		2130				2135					2140						
65	Val	Ser	Phe	Arg	Val	Gly	Leu	His	Glu	Tyr	Pro	Val	Gly	Ser	Gln	Leu	
		2145				2150				2155					2160		

	Pro	Cys	Glu	Pro	Glu	Pro	Asp	Val	Ala	Val	Leu	Thr	Ser	Met	Leu	Thr
				2165						2170					2175	
	Asp	Pro	Ser	His	Ile	Thr	Ala	Glu	Ala	Ala	Gly	Arg	Arg	Leu	Ala	Arg
			2180						2185					2190		
5	Gly	Ser	Pro	Pro	Ser	Met	Ala	Ser	Ser	Ser	Ala	Ser	Gln	Leu	Ser	Ala
		2195					2200						2205			
	Pro	Ser	Leu	Lys	Ala	Thr	Cys	Thr	Ala	Asn	His	Asp	Ser	Pro	Asp	Ala
		2210				2215						2220				
10	Glu	Leu	Ile	Glu	Ala	Asn	Leu	Leu	Trp	Arg	Gln	Glu	Met	Gly	Gly	Asn
	2225				2230					2235					2240	
	Ile	Thr	Arg	Val	Glu	Ser	Glu	Asn	Lys	Val	Val	Ile	Leu	Asp	Ser	Phe
				2245					2250					2255		
	Asp	Pro	Leu	Val	Ala	Glu	Glu	Asp	Glu	Arg	Glu	Val	Ser	Val	Pro	Ala
			2260					2265					2270			
15	Glu	Ile	Leu	Arg	Lys	Ser	Arg	Arg	Phe	Ala	Arg	Ala	Leu	Pro	Val	Trp
		2275					2280					2285				
	Ala	Arg	Pro	Asp	Tyr	Asn	Pro	Pro	Leu	Val	Glu	Thr	Trp	Lys	Lys	Pro
		2290				2295					2300					
20	Asp	Tyr	Glu	Pro	Pro	Val	Val	His	Gly	Cys	Pro	Leu	Pro	Pro	Pro	Arg
	2305				2310					2315					2320	
	Ser	Pro	Pro	Val	Pro	Pro	Pro	Arg	Lys	Lys	Arg	Thr	Val	Val	Leu	Thr
				2325					2330					2335		
	Glu	Ser	Thr	Leu	Ser	Thr	Ala	Leu	Ala	Glu	Leu	Ala	Thr	Lys	Ser	Phe
		2340					2345					2350				
25	Gly	Ser	Ser	Ser	Thr	Ser	Gly	Ile	Thr	Gly	Asp	Asn	Thr	Thr	Thr	Ser
		2355					2360					2365				
	Ser	Glu	Pro	Ala	Pro	Ser	Gly	Cys	Pro	Pro	Asp	Ser	Asp	Val	Glu	Ser
		2370				2375					2380					
30	Tyr	Ser	Ser	Met	Pro	Pro	Leu	Glu	Gly	Glu	Pro	Gly	Asp	Pro	Asp	Leu
	2385				2390					2395					2400	
	Ser	Asp	Gly	Ser	Trp	Ser	Thr	Val	Ser	Ser	Gly	Ala	Asp	Thr	Glu	Asp
				2405					2410				2415			
	Val	Val	Cys	Cys	Ser	Met	Ser	Tyr	Ser	Trp	Thr	Gly	Ala	Leu	Val	Thr
			2420					2425					2430			
35	Pro	Cys	Ala	Ala	Glu	Glu	Gln	Lys	Leu	Pro	Ile	Asn	Ala	Leu	Ser	Asn
		2435					2440					2445				
	Ser	Leu	Leu	Arg	His	His	Asn	Leu	Val	Tyr	Ser	Thr	Thr	Ser	Arg	Ser
		2450				2455					2460					
40	Ala	Cys	Gln	Arg	Gln	Lys	Val	Thr	Phe	Asp	Arg	Leu	Gln	Val	Leu	
	2465				2470				2475				2480			
	Asp	Ser	His	Tyr	Gln	Asp	Val	Leu	Lys	Glu	Val	Lys	Ala	Ala	Ala	Ser
				2485					2490				2495			
	Lys	Val	Lys	Ala	Asn	Leu	Leu	Ser	Val	Glu	Glu	Ala	Cys	Ser	Leu	Thr
			2500					2505				2510				
45	Pro	Pro	His	Ser	Ala	Lys	Ser	Lys	Phe	Gly	Tyr	Gly	Ala	Lys	Asp	Val
		2515					2520					2525				
	Arg	Cys	His	Ala	Arg	Lys	Ala	Val	Ala	His	Ile	Asn	Ser	Val	Trp	Lys
		2530				2535					2540					
50	Asp	Leu	Leu	Glu	Asp	Ser	Val	Thr	Pro	Ile	Asp	Thr	Thr	Ile	Met	Ala
	2545				2550					2555				2560		
	Lys	Asn	Glu	Val	Phe	Cys	Val	Gln	Pro	Glu	Lys	Gly	Gly	Arg	Lys	Pro
				2565				2570					2575			
	Ala	Arg	Leu	Ile	Val	Phe	Pro	Asp	Leu	Gly	Val	Arg	Val	Cys	Glu	Lys
			2580					2585				2590				
55	Met	Ala	Leu	Tyr	Asp	Val	Val	Ser	Lys	Leu	Pro	Leu	Ala	Val	Met	Gly
		2595					2600					2605				
	Ser	Ser	Tyr	Gly	Phe	Gln	Tyr	Ser	Pro	Gly	Gln	Arg	Val	Glu	Phe	Leu
		2610				2615				2620						
60	Val	Gln	Ala	Trp	Lys	Ser	Lys	Lys	Thr	Pro	Met	Gly	Phe	Ser	Tyr	Asp
	2625				2630					2635				2640		
	Thr	Arg	Cys	Phe	Asp	Ser	Thr	Val	Thr	Glu	Ser	Asp	Ile	Arg	Thr	Glu
				2645				2650				2655				
	Glu	Ala	Ile	Tyr	Gln	Cys	Cys	Asp	Leu	Asp	Pro	Gln	Ala	Arg	Val	Ala
			2660				2665					2670				
65	Ile	Lys	Ser	Leu	Thr	Glu	Arg	Leu	Tyr	Val	Gly	Gly	Pro	Leu	Thr	Asn
		2675					2680					2685				

Ser Arg Gly Glu Asn Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly Val
 2690 2695 2700
 Leu Thr Thr Ser Cys Gly Asn Thr Leu Thr Cys Tyr Ile Lys Ala Arg
 2705 2710 2715 2720
 5 Ala Ala Cys Arg Ala Ala Gly Leu Gln Asp Cys Thr Met Leu Val Cys
 2725 2730 2735
 Gly Asp Asp Leu Val Val Ile Cys Glu Ser Ala Gly Val Gln Glu Asp
 2740 2745 2750
 10 Ala Ala Ser Leu Arg Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala
 2755 2760 2765
 Pro Pro Gly Asp Pro Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr
 2770 2775 2780
 Ser Cys Ser Ser Asn Val Ser Val Ala His Asp Gly Ala Gly Lys Arg
 2785 2790 2795 2800
 15 Val Tyr Tyr Leu Thr Arg Asp Pro Thr Thr Pro Leu Ala Arg Ala Ala
 2805 2810 2815
 Trp Glu Thr Ala Arg His Thr Pro Val Asn Ser Trp Leu Gly Asn Ile
 2820 2825 2830
 20 Ile Met Phe Ala Pro Thr Leu Trp Ala Arg Met Ile Leu Met Thr His
 2835 2840 2845
 Phe Phe Ser Val Leu Ile Ala Arg Asp Gln Leu Glu Gln Ala Leu Asn
 2850 2855 2860
 Cys Glu Ile Tyr Gly Ala Cys Tyr Ser Ile Glu Pro Leu Asp Leu Pro
 2865 2870 2875 2880
 25 Pro Ile Ile Gln Arg Leu His Gly Leu Ser Ala Phe Ser Leu His Ser
 2885 2890 2895
 Tyr Ser Pro Gly Glu Ile Asn Arg Val Ala Ala Cys Leu Arg Lys Leu
 2900 2905 2910
 30 Gly Val Pro Pro Leu Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg
 2915 2920 2925
 Ala Arg Leu Leu Ser Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr
 2930 2935 2940
 Leu Phe Asn Trp Ala Val Arg Thr Lys Leu Lys Leu Thr Pro Ile Ala
 2945 2950 2955 2960
 35 Ala Ala Gly Arg Leu Asp Leu Ser Gly Trp Phe Thr Ala Gly Tyr Ser
 2965 2970 2975
 Gly Gly Asp Ile Tyr His Ser Val Ser His Ala Arg Pro Arg Trp Phe
 2980 2985 2990
 40 Trp Phe Cys Leu Leu Leu Leu Ala Ala Gly Val Gly Ile Tyr Leu Leu
 2995 3000 3005
 Pro Asn Arg
 3010

45
 <210> 15
 <211> 21
 <212> DNA
 <213> Artificial Sequence

50
 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

55
 <400> 15
 gaaggaggga ggtttgaagg a

21

60
 <210> 16
 <211> 19
 <212> DNA
 <213> Artificial Sequence

65
 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

<400> 16
ccagttccgg gcaagaact

19

5

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/21002

A. CLASSIFICATION OF SUBJECT MATTER														
IPC(7) : C12N 7/01, 15/63; C12Q 1/70; A61K 39/29														
US CL : 435/5, 6, 235.1, 455, 948; 424/186.1, 199.1, 218.1, 228.1														
According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED														
Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/5, 6, 235.1, 455, 948; 424/186.1, 199.1, 218.1, 228.1														
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched														
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Continuation Sheet														
C. DOCUMENTS CONSIDERED TO BE RELEVANT														
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
Y	WO 00/75337 A1 (THE GOVERNMENT OF THE UNITED STATES OF AMERICA AS REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES) 14 December 2000 (14.12.2000), see entire document.	1-58												
Y	US 2001/0034019 A1 (HONG et al.) 25 October 2001 (25.10.2001), see entire document.	1-58												
Y	LEMON et al. Internal Ribosome Entry Sites Within the RNA Genomes of Hepatitis C Virus and Other Flaviviruses. Seminars in Virology. 1997, Vol. 8, pages 274-288, see entire document.	1-58												
Y	RJINBRAND et al. Mutational Analysis of the GB Virus B Internal Ribosome Entry Site. Journal of Virology. January 2000, Vol. 74, No. 2, pages 773-783, see entire document.	1-58												
Y	FROLOV et al. Cis-Acting RNA Elements Required for Replication of Bovine Viral Diarrhea Virus-Hepatitis C Virus 5' Nontranslated Region Chimeras. RNA. 1998, Vol. 4, pages 1418-1435, see entire document.	1-58												
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"B" earlier application or patent published on or after the international filing date</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"&" document member of the same patent family</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"B" earlier application or patent published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means		"P" document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
"B" earlier application or patent published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family													
"O" document referring to an oral disclosure, use, exhibition or other means														
"P" document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 29 September 2003 (29.09.2003)		Date of mailing of the international search report 28 OCT 2003												
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703)305-3230		Authorized officer <i>Helicia D. Roberts for</i> James L. Grun, Ph.D. Telephone No. 703-308-0196												

INTERNATIONAL SEARCH REPORT

PCT/US03/21002

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ZHAO et al. Poliovirus/Hepatitis C Virus (Internal Ribosome Entry Site-Core) Chimeric Viruses: Improved Growth Properties through Modification of a Proteolytic Cleavage Site and Requirement for Core RNA Sequences but Not for Core-Related Polypeptides. Journal of Virology. February 1999, Vol. 73, No. 2, pages 1546-1554, see entire document.	1-58

INTERNATIONAL SEARCH REPORT

PCT/US03/21002

Continuation of B. FIELDS SEARCHED Item 3:
DIALOG, EAST
search terms: GBV, HCV, chimeric